

Increased risk of mortality associated with social isolation in older men: only when feeling lonely? Results from the Amsterdam Study of the Elderly (AMSTEL)

T. J. Holwerda^{1,2*}, A. T. F. Beekman^{2,3}, D. J. H. Deeg², M. L. Stek^{2,3}, T. G. van Tilburg^{2,4}, P. J. Visser², B. Schmand⁵, C. Jonker² and R. A. Schoevers^{1,2,6}

¹ Department of Psychiatry, ARKIN Institute of Mental Health Care, Amsterdam, The Netherlands

² EMGO Institute, Vrije Universiteit Medisch Centrum Amsterdam (VUMC), Amsterdam Study of the Elderly (AMSTEL), Amsterdam, The Netherlands

³ Department of Psychiatry, INGEEST Institute of Mental Health Care, Amsterdam, The Netherlands

⁴ Department of Social Sciences, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

⁵ Department of Psychology, Universiteit van Amsterdam, Amsterdam, The Netherlands

⁶ Department of Psychiatry, Universitair Medisch Centrum Groningen (UMCG), Groningen, The Netherlands

Background. Loneliness has a significant influence on both physical and mental health. Few studies have investigated the possible associations of loneliness with mortality risk, impact on men and women and whether this impact concerns the situation of being alone (social isolation), experiencing loneliness (feeling lonely) or both. The current study investigated whether social isolation and feelings of loneliness in older men and women were associated with increased mortality risk, controlling for depression and other potentially confounding factors.

Method. In our prospective cohort study of 4004 older persons aged 65–84 years with a 10-year follow-up of mortality data a Cox proportional hazard regression analysis was used to test whether social isolation factors and feelings of loneliness predicted an increased risk of mortality, controlling for psychiatric disorders and medical conditions, cognitive functioning, functional status and sociodemographic factors.

Results. At 10 years follow-up, significantly more men than women with feelings of loneliness at baseline had died. After adjustment for explanatory variables including social isolation, the mortality hazard ratio for feelings of loneliness was 1.30 [95% confidence interval (CI) 1.04–1.63] in men and 1.04 (95% CI 0.90–1.24) in women. No higher risk of mortality was found for social isolation.

Conclusions. Feelings of loneliness rather than social isolation factors were found to be a major risk factor for increasing mortality in older men. Developing a better understanding of the nature of this association may help us to improve quality of life and longevity, especially in older men.

Received 16 December 2010; Revised 13 July 2011; Accepted 4 August 2011; First published online 6 September 2011

Key words: Feelings of loneliness, loneliness, mortality, social isolation.

Introduction

The association between an increased risk of mortality and medical conditions has been extensively investigated. Chronic diseases such as cancer, coronary heart disease, respiratory disease, diabetes mellitus, cerebrovascular disease and dementia are among the most important predictors increasing mortality in older adults (Ostbye *et al.* 1999; Garssen &

Hoogenbeezem, 2005; Donnan *et al.* 2008; Janssen *et al.* 2008). In addition, common mental disorders such as depression substantially increase the risk of death, both by unnatural causes and by their detrimental effects on physical health (Murphy *et al.* 1987; Penninx *et al.* 1999; Geerlings *et al.* 2000; Schoevers *et al.* 2000; Holwerda *et al.* 2007). Much less is known about the association between social conditions such as social isolation and feelings of loneliness and mortality risk.

With age, the number of social relationships decreases. Apart from physical ageing and diminished resilience, older persons are prone to becoming lonely due to loss of intimate relationships. Loneliness has been recognized as a clinically important frailty factor

* Address for correspondence: T. J. Holwerda, M.D., Vrije Universiteit Medisch Centrum Amsterdam, AMTEL-LASA, room A 515, Van der Boerhorststraat 7, 1081 BT, Amsterdam, The Netherlands.

(Email: tj.holwerda@vumc.nl)

in older persons, associated with psychosocial problems, mental disorders such as depression, as well as physical problems. These include increased systolic blood pressure, higher vulnerability for heart conditions, sleeping problems, increased fibrinogen and natural killer cell responses to stress and greater cognitive decline (Cacioppo *et al.* 2002*a,b*; Wilson *et al.* 2003; Steptoe *et al.* 2004; Tilvis *et al.* 2004; Hawkey *et al.* 2006; Wilson *et al.* 2007; Luanaigh & Lawlor, 2008). Evidence emerges that loneliness is at the core of a complex constellation of socio-emotional states that include self-esteem, mood, anxiety, anger, optimism, fear of negative evaluation, shyness, social skills, social support and sociability (Cacioppo *et al.* 2000; Boomsma *et al.* 2005). To date, very few studies have examined whether loneliness may also be associated with longevity, especially when controlling for other potentially confounding factors (Penninx *et al.* 1997).

Loneliness is typically defined in relation to indices of social integration such as marital status, contacts with close friends or family, and lack of social integration and 'embeddedness'. This is often referred to as 'social loneliness' or social isolation (Helmer *et al.* 1999; Fratiglioni *et al.* 2000; De Jong Gierveld & Van Tilburg, 2008; Luanaigh & Lawlor; 2008). However, individuals vary in the way they perceive their social context. Another definition of loneliness is based on the discrepancy between a person's desired and actual relationships. This subjective experience – feeling lonely – is often referred to as perceived or emotional loneliness (De Jong Gierveld & Van Tilburg, 2008; Luanaigh & Lawlor, 2008). Earlier studies generally did not make this distinction between actual contextual factors and the subjective experience of loneliness (Berkman & Syme, 1979; Fratiglioni *et al.* 2000). It is highly relevant to assess whether the untoward consequences of loneliness concern the actual situation of being alone or the perceived lack of embeddedness. It has been argued that networks influence physical and mental health through different pathways such as social support, social influence, access to resources and material goods and through social engagement and attachment. These psychosocial and behavioural processes are thought to influence more proximate pathways to health status including direct physiological stress responses, psychological states and traits including self-esteem, self-efficacy and security, health-damaging and health-promoting behaviour and exposure to diseases. Accordingly, adverse health conditions can be either the result of macrosocial forces, a limited social network and lack of social support (a status of being alone), or a consequence of an emotional state (feelings of loneliness) that relates to biological responses that are potentially

relevant to health, or both (Berkman *et al.* 2000; Steptoe *et al.* 2004). The question remains to what extent social isolation and feelings of loneliness contribute to health outcome.

Apart from being the consequence of (objective) environmental characteristics or the subjective experience of being lonely, loneliness has also been conceptualized as a stable personality trait, with different individual set-points for these feelings in different individuals, and with differences between men and women. Both environmental and genetic components are thought to contribute to these individual variations. Men are thought to be more vulnerable as a result of forming less close and enduring relationships outside marriage (Dykstra & De Jong Gierveld, 2004; Boomsma *et al.* 2005).

Loneliness may also be a symptom of depression. A study by Stek *et al.* (2005), however, showed that people who feel lonely often do not meet the criteria for a depressive disorder. Loneliness and depression are considered to be distinct entities (Luanaigh & Lawlor, 2008).

It thus remains unclear whether social isolation and feelings of loneliness are associated with an increased risk of mortality. Furthermore, to our knowledge no studies have examined whether such an association is different in men and women, although social and individual differences between men and women are present (Berkman *et al.* 2000). Also as far as we know, no studies have been performed to ascertain whether such an association is explained by potentially confounding factors such as depression, somatic illnesses and other health characteristics.

We hypothesized that feelings of loneliness are psychological experiences with adverse effects on health associated with increased mortality, with men being more vulnerable than women. The current study investigates whether feelings of loneliness, rather than social isolation factors, are associated with an increased risk of mortality in men and women in our large cohort of community-dwelling older persons, while controlling for social isolation and potentially confounding influences of mental disorders, somatic conditions and diseases, cognitive functioning, functional status and sociodemographic factors.

Method

Study cohort

Residents of the city of Amsterdam were recruited in The Amsterdam Study of the Elderly (AMSTEL) using the registers of 30 general practitioners (Van Ojen *et al.* 1995; Geerlings *et al.* 2000; Schoevers *et al.* 2000, 2009). Subjects were included if they were over age 65 years

and younger than 84 years, and were randomly selected from four age categories of 5 years each. The population over the age of 65 years (15%) and the age and gender distribution were representative of the Amsterdam population aged 65 years and older. In total, 5666 residents were approached of whom 4051 agreed to participate (71.5%). Non-response in the population aged under 75 years was associated with poor cognitive functioning and with a poor medical condition. In the population aged over 75 years, no associations with non-response were found (Launer *et al.* 1994). A home assessment by means of a structured interview was performed by trained and supervised interviewers. For follow-up analyses mortality data were 98.8% complete, leaving 4004 participants for this study.

Measurements

Assessment of social isolation and feelings of loneliness

At baseline, social isolation and feelings of loneliness were assessed in men and women. Social isolation was operationalized as either living alone or not/no longer being married or by the lack of social support (the question was asked if participants were getting help from family, neighbours or home support). Marital status, living arrangement and social network have earlier been recognized and investigated as risk factors in studies on prognosis of older adults (Helmer *et al.* 1999; Fratiglioni *et al.* 2000). Feelings of loneliness were operationalized as experiencing loneliness (the subjective feeling of loneliness) by the participant when asked by the interviewer: do you feel lonely? Both were assessed by a structured clinical interview (AMSTEL, see below).

Assessment of mortality

The vital status of male and female participants was ascertained up to 1 January 2001 through linkage with the registers of the municipality of Amsterdam or the municipalities in which participants had relocated during the study period. The participants' survival time was calculated on a monthly basis with a follow-up period up to 1 January 2001 with an average follow-up of 117.8 months (109.2 to 127.2 months).

Assessment of potentially confounding and explanatory variables

At baseline, psychiatric disorders (including minor and major depression and generalized anxiety disorder), medical disorders and conditions (including ischaemic heart disease, cerebrovascular disease, respiratory disease, diabetes, hypertension, cancer, arthritis/arthrosis, epilepsy, Parkinson's disease and

dementia), smoking and drinking behaviour, cognitive and daily functioning and demographic characteristics (including educational level) were assessed by means of the structured AMSTEL interview. This structured interview includes the Geriatric Mental State Examination (GMS AGE CAT; Launer *et al.* 1994), a diagnostic system with diagnoses of (neuro) psychiatric disorders (in our study depression, generalized anxiety disorder and dementia) and severity of the disorders on 5 levels. Levels 3–5 represent valid clinical cases – dementia, minor (level 3) and major (level 4, 5) depression, and generalized anxiety disorder – whereas levels 1–2 represent subcases of depression, generalized anxiety disorder and cognitive impairment no dementia (CIND) (Copeland *et al.* 1986, 1987, 1992). The assessment of psychiatric and medical disorders, past and family history on psychiatric and medical disorders and cognitive functioning was performed using the Cambridge Mental Disorders of the Elderly Examination (CAMDEX; Roth *et al.* 1986) which has sections on present and past state of psychiatric and medical functioning and on neuropsychological functioning using the cognitive section of CAMDEX (known as the CAMCOG) and the Mini Mental State Examination (Folstein *et al.* 1975), smoking (20 cigarettes/day during at least 1 year in the past or currently) and drinking (alcohol abuse: four or more/day in the past or currently) behaviour and the activities of daily living (ADL) and instrumental activities of daily living (IADL) scales (Katz *et al.* 1963; Lawton & Brody, 1969).

Statistical analysis

Descriptive (χ^2) statistics were used to characterize baseline data, to assess differences between men and women at baseline on social isolation, feelings of loneliness and potentially confounding factors and to assess whether feeling of loneliness were more prevalent in socially isolated male and female subgroups. Mortality ratios for men and women were calculated for all risk factors potentially associated with excess mortality in bivariate analyses, to assess which risk factors to adjust for in the multivariate model. The associations between the social isolation factors, feelings of loneliness and mortality (when controlling for other variables) were calculated using separate Cox proportional hazard regression models. The interaction between feelings of loneliness and gender was assessed and added to this model. Results are presented as hazard ratios (HRs) for men and women separately, and adjusted for separate social isolation factors and/or feelings of loneliness, depression (also including subthreshold depression), age and education, pre-existing ischaemic heart disease,

Table 1. Characteristics of male and female participants at baseline (n = 4004)^a

Characteristic	All participants, n (%)	Men, n (%)	Women, n (%)	Analysis ^b		
				χ^2	df	p
n	4004	1509	2495			
Age, years				22.8	3	<0.001
65–69	827 (20.7)	347 (23.0)	480 (19.2)			
70–74	959 (24.0)	392 (26.0)	567 (22.7)			
75–79	1039 (25.9)	383 (25.4)	656 (26.3)			
80–86	1179 (29.4)	387 (25.6)	792 (31.7)			
Education				80.7	1	<0.001
>Primary school	2311 (57.7)	1007 (66.7)	1304 (52.3)			
Primary school or less	1693 (42.3)	502 (33.3)	1191 (47.7)			
Social isolation						
Living alone	1827 (45.6)	347 (23.0)	1480 (59.4)	501.1	1	<0.001
Not/no longer married	2049 (51.2)	413 (27.4)	1636 (65.6)	549.8	1	<0.001
Lacking social support	2725 (68.1)	1165 (77.9)	1560 (63.0)	96.1	1	<0.001
Feelings of loneliness	859 (21.5)	180 (12.0)	679 (27.4)	131.3	1	<0.001
Depression	515 (12.9)	105 (7.0)	410 (16.4)	75.3	1	<0.001
Ischaemic heart disease	413 (10.3)	223 (14.8)	190 (7.6)	52.2	1	<0.001
Cerebrovascular disease	229 (5.7)	94 (6.2)	135 (5.4)			n.s.
Diabetes	358 (8.9)	118 (7.8)	240 (9.6)	3.7	1	<0.05
Cancer	444 (11.1)	153 (10.1)	291 (11.7)			n.s.
Respiratory disease	655 (16.4)	305 (20.2)	350 (14.0)	26.3	1	<0.001
Arthritis	684 (17.1)	143 (9.5)	541 (21.7)	98.9	1	<0.001
Epilepsy	69 (1.7)	26 (1.7)	43 (1.7)			n.s.
Parkinson's disease	59 (1.5)	30 (2.0)	29 (1.21)	4.4	1	<0.04
CIND	179 (4.5)	71 (4.7)	108 (4.3)			n.s.
Dementia	258 (6.4)	84 (5.6)	174 (7.0)			n.s.
MMSE				12.1	2	<0.002
26–30	3241 (80.9)	1262 (83.6)	1979 (79.4)			
22–25	517 (12.9)	162 (10.7)	355 (14.2)			
0–21	246 (6.1)	85 (5.6)	161 (6.4)			
ADL disability	320 (8.0)	93 (6.2)	227 (9.1)	10.5	1	<0.001
IADL disability	1022 (25.5)	309 (20.5)	713 (28.6)	32.5	1	<0.001

df, Degrees of freedom; n.s., non-significant ($p > 0.05$); CIND, cognitive impairment no dementia; MMSE, Mini Mental State Examination; ADL, activities of daily living; IADL, instrumental activities of daily living.

^a Data of generalized anxiety disorder, hypertension, cardiac arrhythmias, alcohol abuse and smoking not shown.

^b χ^2 test for differences between men and women in bivariate analysis.

cerebrovascular disease, diabetes, cancer, respiratory disease, arthritis/arthrosis, epilepsy and Parkinson's disease, and cognitive and daily functioning. In order to control for confounding, bivariate associations between depression and social support status were assessed using both a dichotomous and a continuous depression variable incorporating the five severity/caseness levels provided in GMS AGE-CAT. If such an association was found, an interaction term was added to the model in multivariate analyses. Kaplan-Meier curves were created to visualize the association

between feelings of loneliness and excess mortality in men and women.

Results

Characteristics of the study population at baseline (n=4004)

Table 1 summarizes baseline data on the most important variables under review. Men and women differed significantly for most conditions. More

women (27.4%) than men (12.0%) reported feelings of loneliness and more women (59.4%) than men (23.0%) were living alone. Men (72.6%) were more often married than women (34.4%), and women more often were receiving social support (37.0%) than men (22.1%). Male participants had higher education, and more often had ischaemic heart disease, respiratory disease, Parkinson's disease and (a history of/met criteria of) alcohol abuse and smoking. Female participants had higher levels of functional limitations, depression, generalized anxiety disorder, arthritis/arthrosis, hypertension, cardiac arrhythmias and diabetes than men. No differences between men and women were found in diagnosis of epilepsy and cancer, CIND and dementia.

Within subjects who were living alone or who were unmarried, more women than men reported feelings of loneliness ($p < 0.001$). In subjects without social support, more men than women perceived feelings of loneliness ($p < 0.001$) (data not shown).

Feelings of loneliness and associated 10-year mortality (n=4004)

At 10 years follow-up, the overall risk of mortality was 52.3% (men 60.8% and women 47.2%). The mortality rate in persons living alone was 54.4% (men 70.3%, women 50.7%); in those who were not or no longer married this was 55.7% (men 71.9%, women 51.7%). In persons without social support the mortality rate was 45.0% (54.8% of men and 37.6 of women). The mortality rate in older persons with feelings of loneliness was 59.1%; 78.9% of men and 53.9% of women experiencing these feelings had died. Table 2 shows bivariate associations between potential risk factors and 10-year mortality. Living alone, not or no longer being married and feelings of loneliness were associated with an increased risk of mortality in both genders. Mortality ratios were 1.37 [95% confidence interval (CI) 1.19–1.59] for men and 1.29 (95% CI 1.14–1.45) for women living alone, 1.48 (95% CI 1.29–1.70) for men and 1.47 (95% CI 1.30–1.67) for women not or no longer married and 1.71 (95% CI 1.43–2.04) for male participants with feelings of loneliness and 1.31 (95% CI 1.15–1.48) for female participants with these feelings. No increased mortality was found in men and women lacking social support.

Nearly all other potentially confounding factors were also associated with an increased risk of mortality in bivariate analysis, except arthrosis/arthritis, hypertension, generalized anxiety disorder, smoking and alcohol abuse. In participants who did not have social support, an opposite association was found in bivariate analysis. We explained this by hypothesizing that those needing more support were in poorer

health. We found indeed that in participants receiving social support, significantly greater numbers suffered from conditions such as ischaemic heart disease, cerebrovascular disease, respiratory disease, arthrosis/arthritis, diabetes, hypertension, cardiac arrhythmias, Parkinson's disease, depression, generalized anxiety disorder, CIND, dementia, ADL and IADL disabilities (data not shown).

Table 3 shows the association between 10-year risk of mortality and feelings of loneliness in male and female participants, with adjustment for all potentially confounding factors. Results are presented in HRs for time to death. Cox proportional hazard analyses showed that after adjustment for social isolation, age and education, depression (including subthreshold depression), ischaemic heart disease, cerebrovascular disease, diabetes, cancer, respiratory disease, arthritis, epilepsy, Parkinson's disease, cognitive functioning and functional status the association of feelings of loneliness with an increased mortality risk remained significant in men (HR 1.30, 95% CI 1.04–1.62) and lost significance in women (HR 1.01 95% CI 0.87–1.17).

In multivariate analysis no increased risk of mortality was found in socially isolated older adults. HRs were 0.88 (95% CI 0.65–1.19) for men and 0.96 (95% CI 0.78–1.20) for women living alone and 1.21 (95% CI 0.91–1.59) for men and 1.07 (95% CI 0.85–1.34) for women who were not/no longer married (data not shown). However, in men without objective determinants of social isolation (living with others and being married), feelings of loneliness were also associated with an increased risk of mortality. The association between feelings of loneliness and mortality was 1.62 (95% CI 1.14–2.29) for men living with others, HR 1.60 (95% CI 1.13–2.27) for married men, and HR 1.33 (95% CI 0.95–1.87) for men receiving social support. In women living with others, being married or receiving social support, no increased mortality was found in relation to feelings of loneliness (data not shown). Finally, a strong interaction was found between feelings of loneliness and gender ($p < 0.0001$).

In bivariate analysis both the dichotomous and continuous depression variable was not associated with not having social support but instead was associated with receiving social support in both men and women (Pearson χ^2 $p < 0.000$). However, we found no significant interaction between depression and having social support in either men or women ($p = 0.90$ and 0.12) in the multivariate model. Thus, the association between no social support as a measure of social isolation and mortality was not confounded by depression. These data show again that feelings of loneliness independently raise mortality risk and are not dependent on actual social isolation and depression.

Table 2. Bivariate associations between risk factors and excess mortality (10-year follow-up, $n = 4004$)^a

Mortality	All, %	Mortality ratio	Men, %	Mortality ratio	Women, %	Mortality ratio
All	52.3	($n = 2095$)	60.8	($n = 917$)	47.2	($n = 1178$)
Social isolation						
Living alone	50.5	1.09 (1.00–1.19)*	56.1	1.37 (1.19–1.59)*	38.7	1.29 (1.14–1.45)*
(no/yes)	54.4		68.5		48.4	
Marital status	48.7	1.20 (1.10–1.30)*	55.1	1.48 (1.29–1.70)*	36.2	1.47 (1.30–1.67)*
(yes/not or no longer)	55.7		69.6		48.9	
Social support	67.6	0.54 (0.49–0.58)*	78.4	0.47 (0.40–0.54)*	60.2	0.48 (0.43–0.54)*
(yes/lack of)	45.0		53.5		35.7	
Feelings of loneliness	50.2	1.26 (1.14–1.39)*	56.5	1.71 (1.43–2.04)*	42.1	1.31 (1.15–1.48)*
(no/yes)	59.1		77.4		50.9	
Depression	51.1	1.33 (1.18–1.50)*	57.5	2.30 (1.75–2.87)*	43.0	1.30 (1.12–1.50)*
(no/yes)	60.6		80.7		52.3	
Age, years						
65–69 ^b	28.3		37.8		21.1	
70–74	38.5	1.22 (1.12–1.32)*	44.9	1.16 (1.04–1.30)*	31.6	1.29 (1.15–1.46)*
75–79	57.7	2.56 (2.20–2.98)*	68.1	2.56 (2.08–3.17)*	48.4	2.84 (2.28–3.55)*
80–86	75.7	4.18 (3.62–4.83)*	85.6	4.26 (3.47–5.24)*	67.0	4.86 (3.94–6.00)*
Education						
>Primary school	47.6	1.36 (1.25–1.48)*	54.7	1.47 (1.29–1.68)*	39.9	1.45 (1.30–1.63)*
Primary school or less	58.7		68.1		50.0	
Ischaemic heart disease	50.3	1.77 (1.57–2.01)*	57.1	1.42 (1.20–1.68)*	42.7	1.98 (1.65–2.38)*
(no/yes)	70.0		69.7		66.7	
Cerebrovascular disease	51.2	1.76 (1.50–2.06)*	57.8	1.73 (1.36–2.20)*	43.6	1.76 (1.42–2.18)*
(no/yes)	71.6		76.5		62.1	
Cancer	51.5	1.30 (1.14–1.48)*	57.9	1.46 (1.20–1.79)*	43.6	1.26 (1.06–1.49)*
(no/yes)	59.0		68.3		51.5	
Respiratory disease	50.5	1.44 (1.29–1.60)*	56.3	1.53 (1.31–1.78)*	43.3	1.28 (1.09–1.49)*
(no/yes)	61.8		69.6		51.9	
Diabetes	50.6	1.71 (1.50–1.95)*	57.6	1.63 (1.31–2.02)*	42.6	1.84 (1.56–2.18)*
(no/yes)	69.6		75.0		63.1	
Epilepsy	52.0	1.46 (1.04–2.05)*	58.8	1.43 (0.91–2.26)	44.3	1.81 (1.24–2.64)*
(no/yes)	68.1		68.4		58.3	
Parkinson's disease	51.8	2.71 (2.06–3.58)*	58.4	2.55 (1.74–3.75)*	44.0	2.66 (1.79–3.96)*
(no/yes)	88.1		88.9		85.2	
Cognitive impairment	51.6	1.56 (1.30–1.88)*	60.0	1.57 (1.19–2.06)*	46.5	1.55 (1.21–1.98)*
No dementia	67.6		76.1		62.0	
(no/yes)						
Dementia	50.0	2.97 (2.58–3.41)*	58.9	3.27 (2.58–4.14)*	44.5	3.07 (2.58–3.66)*
(no/yes)	86.0		91.7		83.3	
MMSE						
26–30 ^b	47.4		56.5		40.9	
22–25	66.7	1.74 (1.55–1.95)*	74.7	1.90 (1.57–2.31)*	60.2	1.80 (1.55–2.09)*
0–21	87.4	3.40 (2.94–3.92)*	84.4	2.50 (1.70–3.68)*	81.5	3.30 (2.43–4.47)*
ADL disability	50.0	2.49 (2.18–2.84)*	57.2	3.73 (2.98–4.67)*	42.5	2.38 (2.02–2.80)*
(no/yes)	79.4		94.2		68.1	
IADL disability	43.6	2.73 (2.49–2.98)*	53.8	3.02 (2.38–3.83)*	35.4	3.88 (3.23–4.65)*
(no/yes)	77.8		83.1		70.8	

MMSE, Mini Mental State Examination; ADL, activities of daily living; IADL, instrumental activities of daily living.

^a Percentages of mortality in participants without and with condition. Mortality ratios for men and women [deceased or alive on 1 January 2001, % deceased of baseline category, first row mortality (%) without condition, second row with condition unless otherwise specified, odds ratios with 95% confidence intervals]. Data of generalized anxiety disorder, hypertension, cardiac arrhythmias, arthrosis/arthritis, alcohol abuse and smoking are not shown.

^b Reference category.

* Significant mortality ratio ($p < 0.05$).

Table 3. Mortality hazard ratios of feelings of loneliness with time for death, with successive adjustment for potential risk factors and explanatory variables^a

Variable	Mortality hazard ratio (men) of feelings of loneliness	Mortality hazard ratio (women) of feelings of loneliness
Feelings of loneliness	1.71 (1.41–2.07)*	1.28 (1.12–1.46)*
Adjusted for		
Age	1.48 (1.23–1.80)*	1.15 (1.00–1.32)*
Social isolation		
Living alone	1.46 (1.17–1.81)*	1.15 (1.00–1.33)*
Not married	1.46 (1.18–1.81)*	1.15 (1.00–1.32)*
No social support	1.39 (1.12–1.72)*	1.11 (0.96–1.27)
Education	1.38 (1.11–1.72)*	1.10 (0.96–1.27)
Ischaemic heart disease, cerebrovascular disease and diabetes	1.42 (1.14–1.77)*	1.08 (0.94–1.25)
Cancer, respiratory disease, arthritis, epilepsy and Parkinson’s disease	1.39 (1.12–1.73)*	1.08 (0.94–1.25)
Depression	1.31 (1.04–1.64)*	1.07 (0.92–1.24)
Cognitive decline	1.29 (1.03–1.62)*	1.06 (0.92–1.24)
ADL and IADL disability	1.30 (1.04–1.63)*	1.04 (0.90–1.21)

ADL, Activities of daily living; IADL, instrumental activities of daily living.

^a Mortality hazard ratios for men and women (deceased or alive on 1 January 2001) with successive adjustment for potential confounders using Cox proportional hazards model, with 95% confidence intervals.

* Significant mortality ratio ($p < 0.05$).




Fig. 1. Kaplan–Meier survival curves for feelings of loneliness in men and women (proportion of subjects who survived); cumulative survival. (■), Feelings of loneliness; (□), no feelings of loneliness. (a) Association between feelings of loneliness and mortality in men (log rank test 31.8, degrees of freedom 1, $p < 0.000$). (b) Association between feelings of loneliness and mortality in women.

Kaplan–Meier analysis demonstrates that feelings of loneliness are associated with increased mortality in men but not in women (Fig. 1a, b)

Discussion

This study examined whether feelings of loneliness rather than social isolation factors are related to an

increased mortality risk in community-dwelling older persons, and whether this correlation may be different for men and women.

We found that overall, more women than men were socially isolated in terms of not or no longer being married and living alone ($p < 0.001$), but that women were receiving more social support than men ($p < 0.001$). We also found that subjective feelings of

loneliness have a high prevalence in both men (12.0%) and women (27.5%). Although feelings of loneliness were more prevalent in women ($p < 0.001$), such feelings were associated with increased mortality only in older men, when controlling for potentially confounding factors. There was a strong interaction between feelings of loneliness and gender ($p < 0.0001$), confirming a differential association with excess mortality in men and women. After adjustment for social isolation (living alone, not having a marital partner and not receiving social support), depressive disorder (including subthreshold depression), demographic characteristics, common medical disorders, cognitive functioning and functional disabilities, older men who reported feelings of loneliness were 30% more likely to have died at 10-year follow-up. In contrast, social isolation was no longer associated with excess mortality in multivariate analysis.

Interpretation

Feelings of loneliness and excess mortality

Loneliness – like depression, anxiety and anger – has been described as an internal emotional state, with lowered mood and clinical distress negatively influencing mental health and physical well-being. There is some evidence of loneliness stimulating psychobiological pathways involving central nervous system activation of neuroendocrine, autonomic and immune responses with potentially adverse biological effects on physical health (Stepptoe *et al.* 2004). Loneliness may play a causal role in the development and maintenance of depression, but has also been identified as a vulnerability factor that, separate from depression, may contribute to suicidal ideation, suicide attempts, self-inflicted injury and suicide completion (Heinrich & Gullone, 2006).

An explanation for the association between feeling lonely and increased mortality could be that these feelings are an indicator of personality traits such as a low self-esteem, sensitivity to rejection, shyness, deficiency in social skills or a personality disorder (i.e. avoidant and borderline personality disorder). These personality characteristics could accelerate mortality by increasing vulnerability to psychological distress and negative emotions, causing both depression and suppression of anger, factors associated with increased mortality (Wilson *et al.* 2003). Recent findings show that conscientiousness, extraversion and openness are inversely related to all-cause mortality. Persons with these characteristics are more likely to engage in healthy behaviours such as regular exercise, better adherence to medical recommendations, having social interactions that decrease the impact of stress,

and are less likely to be heavy smokers or drinkers (Iwasa *et al.* 2008). Evidence also suggests that personality traits are related to neuroendocrine alterations in hypothalamic–pituitary–adrenal axis function, neuroanatomical alterations in hippocampus and anterior cingulate cortex and alterations in neurotrophin and serotonergic signalling pathways (Foster & MacQueen, 2008). Feelings of loneliness could be concomitant to (one of) these traits, and, like depression, could cause these neuroendocrine and neuroanatomical alterations. The question remains whether the observed increased risk of mortality in older persons with feelings of loneliness is causal or whether these feelings are a (subclinical) manifestation of another condition (e.g. genetic factor, vulnerability/frailty factor or unknown condition).

Gender differences

To our knowledge, previously no hypotheses have been developed to explain the mortality gap between men and women associated with feelings of loneliness. However, from several studies it is known that fundamental biological differences between the genders, such as differences in genetics and the immune system [human leukocyte antigen (HLA) haplotypes] may play a role through a different response to stress between men and women. Oestrogens have a beneficial effect on immune-inflammatory responses, which could also possibly account for the different mortality patterns between men and women (Candore *et al.* 2006; Choi & McLaughlin, 2007; Oksuzyan *et al.* 2008). From studies on loneliness, depression, health-seeking behaviour and longevity, we know that older men are particularly reluctant to disclose emotional distress and are more likely to find an intimate attachment in marriage, whereas women may also find protection from loneliness in other social ties. For men, more than women, a marriage bond appears to play a pivotal role in the involvement with others (Van Grootheest *et al.* 1999; Murray *et al.* 2006). With men possibly being more vulnerable to emotional distress and to loss of an intimate marriage attachment, this could account for the mortality gap in feelings of loneliness.

Strengths and limitations

The strengths of this study include its large sample size, with a virtually complete ascertainment of vital status over a long (10-year) period. Furthermore, the study contains a wide range of medical, individual and social risk factors that, in this combination, are rarely available in other studies. The fact that the study was performed in community-dwelling older persons underlines the importance of these findings

for public health. Data were collected by means of a structured interview performed by trained and supervised interviewers. Patients' self-report on chronic diseases and conditions have been shown to be fairly accurate (Kriegsman *et al.* 1996). By analysing the effects of both an objective estimation of social embeddedness and the individual perception of loneliness, the study was able to focus on the actual meaning that people attach to such circumstances, in line with a deepening understanding of the concept of loneliness.

Some limitations also warrant discussion. First, despite the ability to distinguish between feelings of loneliness and social isolation, we were not able to use a more extensive loneliness scale for measuring the severity and extent of loneliness and thereby ascertain structural aspects of social relationships (such as number of contacts and network size). At the time of the first wave of the study, the questionnaire used represented a reliable instrument for measuring social integration, and there are obvious advantages to using a short questionnaire when interviewing large numbers of respondents. However, this simplicity is also a weakness. The question 'Do you feel lonely?' presumes an understanding of the concept of loneliness by participants. Both the nature and meaning of the concept may vary among different groups of people and over time. Still, the use of a single-question self-rating scale to measure loneliness has been widely used in both Europe and North America. It was found to be highly acceptable to research participants. By asking directly about loneliness, it offers the possibility of describing a personal experience, while loneliness scales ask about loneliness in an indirect way by relating it to social networks and the availability of relationships. Especially in older persons, such a single question may not necessarily be less valid as older persons are less likely to admit feeling lonely as they consider loneliness as stigmatizing (Victor, 2005). In the primary-care setting where time is often limited, the use of a limited number of questions is time efficient and mandatory and does appear to identify people with a problem of clinical significance that needs to be addressed by health-care workers (Luanaigh & Lawlor, 2008). A second potential limitation is that our definition of social isolation might be vulnerable to bias by depression. Reports about living alone and not being married are straightforward and probably not affected by depression status, but depressed persons may have an unduly negative view of their relationship to their social environment (Beck *et al.* 1961) and complaints of loneliness in elderly depressed patients may overemphasize actual isolation (Beck *et al.* 1961; Kay *et al.* 1964; Morgado *et al.* 1991a,b). We did find that depression was more

prevalent in older persons who were receiving social support. However, we found no interaction between depression and social support in the multivariate model. A third limitation is that we are unable to exclude the possibility of residual confounding due to errors in the measurement of confounding variables or from measures which were not assessed. Examples of these are personality and personality disorders. Last, as our study was performed in a Caucasian population we do not know if our findings are generalizable to non-Caucasian populations.

Conclusions, implications for future research and therapeutic interventions

In later life, feelings of loneliness significantly increase the risk of death in older men, also when taking into account a comprehensive set of potentially confounding other risk factors associated with longevity. As this association remained after controlling for different aspects of social isolation, feeling lonely seems to have more to do with individual perceptions of (the quality of) social contacts and interactions than with the objective situation of being alone. Feelings of loneliness indicate a warning signal for approaching death in older men. Future work will need to establish whether feelings of loneliness are an indicator of central nervous system activation of neuroendocrine, autonomic and immune responses with adverse biological effects, an indicator of vulnerable personality traits or other unknown conditions accelerating mortality. It will also need to focus on the somatic and psychosocial differences between men and women with respect to these feelings. As an example, men who feel lonely may be more prone to develop an unhealthy life-style, or self-injurious and suicidal behaviour. Future research on loneliness using more extensive measures should seek to disentangle the different pathways involved in loneliness such as macrosocial forces, social networks, social support and emotional states and their relation to behavioural and biological responses including higher morbidity and mortality.

In terms of intervention, a number of approaches have been suggested to address social isolation. These include general education and information to inform the general public about the problem, promotion of expertise on social isolation in health-care professionals, social and personal activation of socially isolated individuals, courses to improve social skills and psychotherapy, e.g. dealing with the loss of a partner (Cattan *et al.* 2005; Van Tilburg & De Jong Gierveld, 2007). Further study is needed to clarify whether these interventions also have an impact on feeling lonely, and whether interventions would also be effective in reducing the excess risk of mortality in men.

Feelings of loneliness are a considerable individual risk, promoting distress and adverse health consequences that deserve clinical attention. Developing a better understanding of the nature of these feelings may help us to improve care and possibly develop interventions that may both enhance quality of life and increase lifespan, especially in older men.

Acknowledgements

For funding, this study received grants from The Netherlands Health Research Promotion Programme and The Netherlands Foundation of Mental Health. The study was approved by the Medical Ethical Committee, Vrije Universiteit Amsterdam.

Declaration of Interest

None.

References

- Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J (1961). An inventory for measuring depression. *Archives of General Psychiatry* **4**, 561–571.
- Berkman LF, Glass T, Brisette I, Seeman TE (2000). From social integration to health: Durkheim in the new millennium. *Social Science and Medicine* **51**, 843–857.
- Berkman LF, Syme SL (1979). Social networks, host resistance, and mortality: a nine-year follow-up study of Alameda county residents. *American Journal of Epidemiology* **109**, 186–204.
- Boomsma DI, Willemsen G, Dolan CV, Hawky LC, Cacioppo JT (2005). Genetic and environmental contributions to loneliness in adults: The Netherlands Twin Register Study. *Behavior Genetics* **35**, 745–752.
- Cacioppo JT, Ernst JM, Bursleson MH, McClintock MK, Malarkey WB, Kowalewski RB, Paulsen A, Hobson JA, Hugdahl K, Spiegel D, Berntson GG (2000). Lonely traits and concomitant physiological processes: the MacArthur social neuroscience studies. *International Journal of Psychophysiology* **35**, 143–154.
- Cacioppo JT, Hawkey LC, Berntson GG, Ernst JM, Gibbs AC, Stickgold R, Hobson JA (2002a). Do lonely days invade the nights? Potential social modulation of sleep efficiency. *Psychological Science* **13**, 384–387.
- Cacioppo JT, Hawkey LC, Crawford E, Ernst JM, Bursleson MH, Kowalewski RB, Malarkey WB, Cauter EV, Berntson GG (2002b). Loneliness and health: potential mechanisms. *Psychosomatic Medicine* **64**, 407–417.
- Candore G, Balisteri CR, Listi F, Grimaldi MP, Vasto S, Colonna-Romano G, Franceschi C, Lio D, Caselli G, Caruso C (2006). Immunogenetics, gender and longevity. *Annals of the New York Academy of Sciences* **1089**, 516–537.
- Cattan M, White M, Bond J, Learmouth A (2005). Preventing social isolation and loneliness among older people: a systematic review of health promotion interventions. *Ageing and Society* **25**, 41–67.
- Choi BG, McLaughlin MA (2007). Why men's hearts break: cardiovascular effects of sex steroids. *Endocrinology Metabolism Clinics of North America* **36**, 365–377.
- Copeland JR, Davidson IA, Dewey ME, Gilmore C, Larkin BA, McWilliam C, Saunders PA, Scott A, Sharma V, Sullivan C (1992). Alzheimer's disease, other dementias, depression and pseudodementia: prevalence, incidence and three-year outcome in Liverpool. *British Journal of Psychiatry* **161**, 230–239.
- Copeland JR, Dewey ME, Giffith-Jones HM (1986). A computerized psychiatric diagnostic system and case nomenclature for elderly subjects: GMS AGE CAT. *Psychological Medicine* **16**, 89–99.
- Copeland JR, Dewey ME, Wood N, Searle R, Davidson IA, McWilliam C (1987). Range of mental illness among the elderly in the community. Prevalence in Liverpool using the GMS-AGE CAT package. *British Journal of Psychiatry* **150**, 815–823.
- De Jong Gierveld J, Van Tilburg T (2008). A shortened scale for overall, emotional and social loneliness. *Tijdschrift voor Gerontologie en Geriatrie* **39**, 4–15.
- Donnan GA, Fisher M, Macleod M, Davis SM (2008). Stroke. *Lancet* **371**, 1612–1623.
- Dykstra PA, De Jong Gierveld J (2004). Gender and marital-history differences in emotional and social loneliness among Dutch older adults. *Canadian Journal on Aging* **23**, 141–155.
- Folstein MF, Folstein SF, McHugh PR (1975). 'Mini mental state'. A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research* **12**, 189–198.
- Foster JA, MacQueen G (2008). Neurobiological factors linking personality traits and major depression. *Canadian Journal of Psychiatry* **53**, 6–13.
- Fratiglioni L, Wang HX, Ericsson K, Maytan M, Winblad B (2000). Influence of social network on occurrence of dementia: a community-based longitudinal study. *Lancet* **355**, 1315–1319.
- Garssen MJ, Hoogenbeezem J (2005). Backgrounds on recent trends in mortality in The Netherlands. *Nederlands Tijdschrift voor Geneeskunde* **149**, 2554–2560.
- Geerlings MI, Schoevers RA, Beekman ATF, Jonker C, Deeg DJH, Schmand B, Ader HJ, Bouter LM, Van Tilburg W (2000). Depression and risk of cognitive decline and Alzheimer's disease. Results of two prospective community based studies in the Netherlands. *British Journal of Psychiatry* **176**, 568–575.
- Hawkey LC, Masi CM, Berry JD, Cacioppo JT (2006). Loneliness is a unique predictor of age-related differences in systolic blood pressure. *Psychology and Aging* **21**, 152–164.
- Heinrich LM, Gullone E (2006). The clinical significance of loneliness: a literature review. *Clinical Psychology Review* **26**, 695–718.
- Helmer C, Damon D, Letenneur L, Fabrigoule C, Barberger-Gateau P, Lafont S, Fuhrer R, Antonucci T, Commenges D, Orgogozo M, Dartigues JF (1999). Marital status and risk of Alzheimer's disease. A French population-based cohort study. *Neurology* **53**, 1953–1958.

- Holwerda TJ, Schoevers RA, Dekker J, Deeg DJH, Jonker C, Beekman ATF (2007). The relationship between generalized anxiety disorder, depression and mortality in old age. *International Journal of Geriatric Psychiatry* **22**, 241–249.
- Iwasa H, Masui Y, Gondo Y, Inagaki H, Kawaai C, Suzuki T (2008). Personality and all-cause mortality among older adults dwelling in a Japanese community: a five-year population-based prospective cohort study. *American Journal of Geriatric Psychiatry* **16**, 399–405.
- Janssen F, Kunst AE, Mackenbach JP (2008). Variations in the pace of mortality in elderly in 7 north-western European countries between 1950–1999: the impact of smoking. *Nederlands Tijdschrift voor Geneeskunde* **152**, 1478–1484.
- Katz S, Ford AB, Moskowitz RW, Jackson BA, Jaffe MW (1963). Studies of illness in the aged. *The index of ADL: a standardized measure of biological and psychosocial function. Journal of the American Medical Association* **185**, 914–919.
- Kay DWK, Beamish P, Roth M (1964). Old age mental disorders in Newcastle upon Tyne: a study of prevalence. *British Journal of Psychiatry* **110**, 146–158.
- Kriegsman DMW, Penninx BWJH, Van Eijk JThM, Boeke AJP, Deeg DJH (1996). Self-reports and general practitioner information on the presence of chronic diseases in community dwelling elderly. A study on the accuracy of patients' self reports and on determinants of inaccuracy. *Journal of Clinical Epidemiology* **49**, 1407–1417.
- Launer LJ, Wind AW, Deeg DJ (1994). Nonresponse pattern and bias in a community-based cross-sectional study of cognitive functioning among the elderly. *American Journal of Epidemiology* **139**, 803–812.
- Lawton MP, Brody EM (1969). Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist* **9**, 179–186.
- Lunaigh CO, Lawlor BA (2008). Loneliness and the health of older people. *International Journal of Geriatric Psychiatry* **23**, 1213–1221.
- Morgado A, Raoux N, Jourdain G, Lecrubier Y, Widlocher D (1991a). Over-reporting of maladjustment by depressed subjects. Findings from retesting after recovery. *Social Psychiatry and Psychiatric Epidemiology* **26**, 68–74.
- Morgado A, Smith M, Lecrubier Y, Widlocher D (1991b). Depressed subjects unwittingly overreport poor social adjustment which they reappraise when recovered. *Journal of Nervous and Mental Disease* **179**, 614–619.
- Murphy JM, Monson RR, Olivier DC (1987). Affective disorders and mortality. A general population based study. *Archives of General Psychiatry* **44**, 473–480.
- Murray J, Banerjee S, Byng R, Tylee A, Bhugra D, McDonald A (2006). Primary care professionals' perceptions of depression in older people: a qualitative study. *Social Science and Medicine* **63**, 1363–1373.
- Oksuzyan A, Juel K, Vaupel JW, Christensen K (2008). Men: good health and high mortality. Sex differences in health and aging. *Aging Clinical Experimental Research* **20**, 91–102.
- Ostbye T, Hill G, Steenhuis R (1999). Mortality in elderly Canadians with and without dementia. A 5-year follow-up. *Neurology* **53**, 521–532.
- Penninx BWJH, Geerlings SW, Deeg DJH, Van Eijk JT, van Tilburg W, Beekman AT (1999). Minor and major depression and the risk of death in older persons. *Archives of General Psychiatry* **56**, 889–895.
- Penninx BWJH, Van Tilburg T, Kriegsman DMW, Deeg DJH, Boeke AJP, Van Eijk JTM (1997). Effects of social support and personal coping resources on mortality in older age: The Longitudinal Aging Study Amsterdam. *American Journal of Epidemiology* **146**, 510–519.
- Roth M, Tym E, Mountjoy CQ, Huppert FA, Hendrie H, Verma S, Goddard R (1986). CAMDEX. A standardised instrument for the diagnosis of mental disorder in the elderly with special reference to the early detection of dementia. *British Journal of Psychiatry* **149**, 698–709.
- Schoevers RA, Geerlings MI, Beekman ATF, Penninx BWJH, Deeg DJH, Jonker C, Van Tilburg W (2000). The association of depression and gender with mortality in late life. *British Journal of Psychiatry* **177**, 336–342.
- Schoevers RA, Geerlings MI, Deeg DJ, Holwerda TJ, Jonker C, Beekman AT (2009). Depression and excess mortality: evidence for a dose response relation in community living elderly. *International Journal of Psychiatry* **24**, 169–176.
- Stek ML, Vinkers DJ, Gussekloo J, Beekman ATF, Van der Mast RC, Westendorp RGJ (2005). Is depression in old age fatal only when people feel lonely? *American Journal of Psychiatry* **162**, 178–180.
- Stephoe A, Owen N, Kunz-Ebrecht SR, Brydon L (2004). Loneliness and neuroendocrine, cardiovascular and inflammatory stress responses in middle-aged men and women. *Psychoneuroendocrinology* **29**, 593–611.
- Tilvis RS, Kähönene-Väre MH, Jolkonen J, Valvanne J, Pitkala KH, Strandberg TE (2004). Predictors of cognitive decline and mortality of aged people over a 10-year period. *Journal of Gerontology* **59**, 268–274.
- Van Grootheest DS, Beekman ATF, Broese van Groenou MI, Deeg DJH (1999). Sex differences in depression after widowhood. Do men suffer more? *Social Psychiatry and Psychiatric Epidemiology* **34**, 391–398.
- Van Ojen R, Hooijer C, Jonker C, Lindeboom J, Van Tilburg W (1995). Late-life depressive disorder in the community, early onset and the decrease of vulnerability with increasing age. *Journal of Affective Disorders* **33**, 159–166.
- Van Tilburg T, De Jong Gierveld (2007). *Zicht op eenzaamheid. Achtergronden, oorzaken en aanpak (Sight on Loneliness. Backgrounds, Origins and Approach)*. Van Gorcum: Assen, The Netherlands.
- Victor C (2005). Measuring loneliness in later life: a comparison of differing measures. *Reviews in Clinical Gerontology* **15**, 63–70.
- Wilson RS, Bienias JL, Mendes de Leon CF, Evans DA, Bennett DA (2003). Negative affect and mortality in older persons. *American Journal of Epidemiology* **158**, 827–835.
- Wilson RS, Krueger KR, Arnold SE, Schneider JA, Kelly JF, Barnes LL, Tang Y, Bennett DA (2007). Loneliness and risk of Alzheimer disease. *Archives of General Psychiatry* **64**, 234–240.