

TABLE OF CONTENTS

Chapter 1	General Introduction Adapted from: Conventional versus biocompatible peritoneal dialysis fluids: more questions than answers? <i>Nephrology Dialysis Transplantation Plus. 2008;1(Suppl 4):iv46-iv50.</i>	9
Chapter 2	Initiation of peritoneal dialysis in the first weeks after catheter insertion: a comparison of a neutral pH, low GDP PD fluid and a conventional PD fluid <i>Clinical Nephrology 2018;89(2):75-82.</i>	29
Chapter 3	The first weeks on peritoneal dialysis: the effects on clinical parameters <i>Advances in Peritoneal Dialysis. 2007;23:102-6.</i>	43
Chapter 4	Effects of conversion to a Bicarbonate/Lactate-Buffered, neutral-pH, low-GDP PD regimen in prevalent PD: a 2-year randomized clinical trial <i>Peritoneal Dialysis International 2017;37(3):273-282.</i>	53
Chapter 5	Pharmacological targets and peritoneal membrane remodeling <i>Peritoneal Dialysis International 2014;34(1):114-23.</i>	73
Chapter 6	Protective effects of paricalcitol on peritoneal remodeling during peritoneal dialysis <i>Biomed Research International 2015;2015:468574.</i>	93
Chapter 7	Effects of oral paricalcitol and calcitriol treatment on peritoneal membrane characteristics of peritoneal dialysis patients; a pilot study <i>Peritoneal Dialysis International 2018 Jan 31. pii: pdi.2017.00150. [Epub ahead of print]</i>	115
Chapter 8	Is vitamin D lost in peritoneal dialysis effluent? <i>Submitted</i>	133
Chapter 9	Summary, General discussion and Future perspective	147
Addendum	Nederlandse samenvatting voor niet ingewijden	159
	Dankwoord	167