

References

- [1] Abegaz, F. and Wit, E. (2013). Sparse time series chain graphical models for reconstructing genetic networks. *Biostatistics*, 1:1–14.
- [2] Abramovich, F. and Angelini, C. (2006). Bayesian Maximum a posteriori Multiple Testing Procedure. *The Indian Journal of Statistics*, 68:436–460.
- [3] Agarwal, V., Bell, G. W., Nam, J. W., and Bartel, D. P. (2015). Predicting effective microRNA target sites in mammalian mRNAs. *eLife*, 4:e05005.
- [4] Alix-Panabières, C. and Pantel, K. (2013). Circulating tumor cells: liquid biopsy of cancer. *Clinical Chemistry*, 59(1):110–118.
- [5] Allen, G. I. and Liu, Z. (2013). A local Poisson graphical model for inferring networks from sequencing data. *IEEE Transactions on NanoBioscience*, 12(3):189–198.
- [6] Alon, U. (2003). Biological networks: the tinkerer as an engineer. *Science*, 301:1866–1867.
- [7] Alon, U. (2007). Network motifs: theory and experimental approaches. *Nature Reviews Genetics*, 8(6):450–461.
- [8] Angelini, C., De Canditiis, D., Mutarelli, M., and Pensky, M. (2007). A Bayesian approach to estimation and testing in time-course microarray experiments. *Statistical Applications in Genetics and Molecular Biology*, 6:1–30.
- [9] Anwar, S. L., Albat, C., Krech, T., Hasemeier, B., Schipper, E., Schweitzer, N., Vogel, A., Kreipe, H., and Lehmann, U. (2013). Concordant hypermethylation of intergenic microRNA genes in human hepatocellular carcinoma as new diagnostic and prognostic marker. *International Journal of Cancer*, 133(3):660–670.
- [10] Balaguer, F., Link, A., Lozano, J. J., Cuatrecasas, M., Nagasaka, T., Boland, C. R., and Goel, A. (2010). Epigenetic silencing of miR-137 is an early event in colorectal carcinogenesis. *Cancer Research*, 70(16):6609–6618.
- [11] Band, V., Zajchowski, D., Kulesa, V., and Sager, R. (1990). Human papilloma virus DNAs immortalize normal human mammary epithelial cells and reduce their growth factor requirements. *Proceedings of the National Academy of Sciences*, 87(1):463–467.
- [12] Bandres, E., Agirre, X., Bitarte, N., Ramirez, N., Zarate, R., Roman-Gomez, J., Prosper, F., and Garcia-Foncillas, J. (2009). Epigenetic regulation of microRNA

- expression in colorectal cancer. *International Journal of Cancer*, 125(11):2737–2743.
- [13] Bar-Joseph, Z., Gerber, G. K., Lee, T. I., Rinaldi, N. J., Yoo, J. Y., Robert, F., Gordon, D. B., Fraenkel, E., Jaakkola, T. S., Young, R. A., and Gifford, D. K. (2003). Computational discovery of gene modules and regulatory networks. *Nature Biotechnology*, 21:1337–1324.
- [14] Baxter, R. C. (2003). Insulin-like growth factor (IGF)-binding proteins: interactions with IGFs and intrinsic bioactivities. *American Journal of Physiology-Endocrinology and Metabolism*, 278:967–976.
- [15] Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)*, 57:289–300.
- [16] Bickel, P. J. and Doksum, K. A. (2001). *Mathematical Statistics*. Prentice Hall, New Jersey.
- [17] Bier, A., Giladi, N., Kronfeld, N., Lee, H. K., Cazacu, S., Finniss, S., Xiang, C., Poisson, L., et al. (2013). MicroRNA-137 is downregulated in glioblastoma and inhibits the stemness of glioma stem cells by targeting RTVP-1. *Oncotarget*, 4(5):665.
- [18] Bierkens, M., Krijgsman, O., Wilting, S. M., Bosch, L., Jaspers, A., Meijer, G. A., Meijer, C. J. L. M., Snijders, P. J. F., Ylstra, B., and Steenbergen, R. D. M. (2013). Focal aberrations indicate EYA2 and hsa-miR-375 as oncogene and tumor suppressor in cervical carcinogenesis. *Genes, Chromosomes and Cancer*, 52:56–68.
- [19] Bilgrau, A. E., Peeters, C. F. W., Eriksen, P. S., Bøgsted, M., and Van Wieringen, W. N. (2015). Targeted fused ridge estimation of inverse covariance matrices from multiple high-dimensional data classes. <http://arxiv.org/pdf/1509.07982.pdf>, 1:1.
- [20] Boldstad, B. M., Irizarry, R. A., Astrand, M., and Speed, T. P. (2003). A comparison of normalization methods for high density oligonucleotide array data based on bias and variance. *Bioinformatics*, 19(2):185–193.
- [21] Botezatu, A., Goia-Rusanu, C. D., Iancu, I. V., Huica, I., Plesa, A., Socolov, D., Ungureanu, C., and Anton, G. (2011). Quantitative analysis of the relationship between microRNA-124a,-34b and-203 gene methylation and cervical oncogenesis. *Molecular Medicine Reports*, 4:121–8.
- [22] Brenet, F., Moh, M., Funk, P., Feierstein, E., Viale, A. J., Socci, N. D., and Scandura, J. M. (2011). DNA methylation of the first exon is tightly linked to transcriptional silencing. *PloS One*, 6(1):e14524.
- [23] Buffart, T. E., Overmeer, R. M., Steenbergen, R. D. M., Tijssen, M., van Grieken, N. C. T., Snijders, P. J. F., Grabsch, H. I., van de Velde, C. J. H., Carvalho, B., and Meijer, G. A. (2008). MAL promoter hypermethylation as a novel prognostic marker in gastric cancer. *British Journal of Cancer*, 99(11):1802.

- [24] Buttitta, F., Pellegrini, C., Marchetti, A., Gadducci, A., Cosio, S., Felicioni, L., Barassi, F., Salvatore, S., Martella, C., Coggi, G., et al. (2003). Human telomerase reverse transcriptase mRNA expression assessed by real-time reverse transcription polymerase chain reaction predicts chemosensitivity in patients with ovarian carcinoma. *Journal of Clinical Oncology*, 21(7):1320–1325.
- [25] Charbonnier, C., Chiquet, J., and Ambroise, C. (2010). Weighted-LASSO for structured network inference from time course data. *Statistical Applications in Genetics and Molecular Biology*, 9:1–27.
- [26] Chen, C., Ridzon, D. A., Broomer, A. J., Zhou, Z., Lee, D. H., Nguyen, J. T., Barbisin, M., Xu, N. L., Mahuvakar, V. R., Andersen, M. R., et al. (2005). Real-time quantification of microRNAs by stem-loop RT-PCR. *Nucleic Acids Research*, 33(20):e179–e179.
- [27] Chen, Q., Chen, X., Zhang, M., Fan, Q., Luo, S., and Cao, X. (2011). miR-137 is frequently down-regulated in gastric cancer and is a negative regulator of Cdc42. *Digestive Diseases and Sciences*, 56(7):2009–2016.
- [28] Chen, T. M., Pecoraro, G., and Defendi, V. (1993). Genetic analysis of in vitro progression of human papillomavirus-transfected human cervical cells. *Cancer Research*, 53:1167–1171.
- [29] Chen, X., Hu, H., Guan, X., Xiong, G., Wang, Y., Wang, K., Li, J., Xu, X., Yang, K., and Bai, Y. (2012). CpG island methylation status of miRNAs in esophageal squamous cell carcinoma. *International Journal of Cancer*, 130(7):1607–1613.
- [30] Chen, X., Ruan, A., Wang, X., Han, W., Wang, R., Lou, N., Ruan, H., Qiu, B., Yang, H., and Zhang, X. (2014). miR-129-3p, as a diagnostic and prognostic biomarker for renal cell carcinoma, attenuates cell migration and invasion via downregulating multiple metastasis-related genes. *Journal of Cancer Research and Clinical Oncology*, 140(8):1295–1304.
- [31] Chen, X., Zhang, L., Zhang, T., Hao, M., Zhang, X., Zhang, J., Xie, Q., Wang, Y., Guo, M., Zhuang, H., et al. (2013). Methylation-mediated repression of microRNA 129-2 enhances oncogenic SOX4 expression in HCC. *Liver International*, 33(3):476–486.
- [32] Chow, L. T., Broker, T. R., and Steinberg, B. M. (2010). The natural history of human papillomavirus infections of the mucosal epithelia. *Apmis*, 118(6-7):422–449.
- [33] Colak, S. and ten Dijke, P. (2017). Targeting TGF- β signaling in cancer. *Trends in Cancer*.
- [34] Cover, T. M. and Thomas, J. A. (2006). *Elements of Information Theory*. John Wiley, New York.
- [35] Crainiceanu, C. M., Ruppert, D., and Wand, M. P. (2005). Bayesian Analysis for Penalized Spline Regression using WinBUGS. *Journal of Statistical Software*, 14:1–47.

- [36] Dahl, J., Roychowdhury, V., and Vandenberghe, L. (2005). Maximum likelihood estimation of Gaussian graphical models: numerical implementation and topology selection. *Technical Report*, 1:1.
- [37] Dahlhaus, R. (2000). Graphical interaction models for multivariate time series. *Metrika*, 51(2):157–172.
- [38] Dahlhaus, R. and Eichler, M. (2003). Causality and graphical models in time series analysis. *Oxford Statistical Science Series*, 1:115–137.
- [39] Den Boon, J. A., Pyeon, D., Wang, S. S., Horswill, M., Schiffman, M., Sherman, M., Zuna, R. E., Wang, Z., Hewitt, S. M., Pearson, R., et al. (2015). Molecular transitions from papillomavirus infection to cervical precancer and cancer: Role of stromal estrogen receptor signaling. *Proceedings of the National Academy of Sciences*, 112(25):3255–3264.
- [40] D’Ercole, A. J. and Calikoglu, A. S. (2001). The case of local versus endocrine IGF-I actions: the jury is still out. *Growth Hormone & IGF Research*, 11:261–265.
- [41] Dobra, A., Lenkoski, A., et al. (2011). Copula Gaussian graphical models and their application to modeling functional disability data. *The Annals of Applied Statistics*, 5(2A):969–993.
- [42] Doorbar, J. (2006). Molecular biology of human papillomavirus infection and cervical cancer. *Clinical Science*, 110:525–541.
- [43] Du, Y., Liu, Z., Gu, L., Zhou, J., Zhu, B.-d., Ji, J., and Deng, D. (2012). Characterization of human gastric carcinoma-related methylation of 9 miR CpG islands and repression of their expressions in vitro and in vivo. *BMC Cancer*, 12(1):1.
- [44] Duensing, S. and Münger, K. (2002). Human papillomaviruses and centrosome duplication errors: modeling the origins of genomic instability. *Oncogene*, 21(40):6241.
- [45] Duensing, S. and Münger, K. (2004). Mechanisms of genomic instability in human cancer: insights from studies with human papillomavirus oncoproteins. *International Journal of Cancer*, 109:157–162.
- [46] Dürst, M., Dzarlieva-Petrusevska, R., Boukamp, P., Fusenig, N., and Gissmann, L. (1987). Molecular and cytogenetic analysis of immortalized human primary keratinocytes obtained after transfection with human papillomavirus type 16 DNA. *Oncogene*, 1(3):251–256.
- [47] Eads, C. A., Danenberg, K. D., Kawakami, K., Saltz, L. B., Blake, C., Shibata, D., Danenberg, P. V., and Laird, P. W. (2000). MethyLight: a high-throughput assay to measure DNA methylation. *Nucleic Acids Research*, 28(8):e32–00.
- [48] Eckel, J. E., Gennings, C., Chinchilli, V. M., Burgoon, L. D., and Zacharewski, T. R. (2004). Empirical Bayes gene screening tool for time-course or dose-response microarray data. *Journal of Biopharmaceutical Statistics*, 14:647–670.

- [49] Efron, B. (2004). Large-scale simultaneous hypothesis testing: the choice of a null hypothesis. *Journal of the American Statistical Association*, 99(465):96–104.
- [50] Efron, B., Tibshirani, R., Storey, J. D., and Tusher, V. (2001). Empirical Bayes analysis of a microarray experiment. *Journal of the American Statistical Association*, 96:1151–1160.
- [51] Ernst, J., Nau, G. J., and Bar-Joseph, Z. (2005). Clustering short time series gene expression data. *Bioinformatics*, 21:i159–i168.
- [52] Evans, A. J., Russell, R. C., Roche, O., Burry, T. N., Fish, J. E., Chow, V. W. K., Kim, W. Y., Saravanan, A., Maynard, M. A., Gervais, M. L., et al. (2007). VHL promotes E2 box-dependent E-cadherin transcription by HIF-mediated regulation of SIP1 and snail. *Molecular and Cellular Biology*, 27(1):157–169.
- [53] Faith, J. J., Hayete, B., Thaden, J. T., Mogno, I., Wierzbowski, J., Cottarel, G., Kasif, S., Collins, J. J., and Gardner, T. S. (2007). Large-scale mapping and validation of escherichia coli transcriptional regulation from a compendium of expression profiles. *PLoS Biology*, 5(1):e8.
- [54] Fan, D. M., Tian, X. Y., Wang, R. F., and Yu, J. J. (2014). The prognosis significance of TGF- β 1 and ER protein in cervical adenocarcinoma patients with stage Ib~IIa. *Tumor Biology*, 35(11):11237–11242.
- [55] Fan, J. and Li, R. (2001). Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of the American Statistical Association*, 96:1348–1360.
- [56] Ferlay, J., Soerjomataram, I., Dikshit, R., Eser, S., Mathers, C., Rebelo, M., Parkin, D., Forman, D., and Bray, F. (2015). Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *International Journal of Cancer*, 136:359–386.
- [57] Freedman, V. H. and Shin, S. (1974). Cellular tumorigenicity in nude mice: correlation with cell growth in semi-solid medium. *Cell*, 3:355–359.
- [58] Friedman, J., Hastie, T., and Tibshirani, R. (2008). Sparse inverse covariance estimation with the graphical lasso. *Biostatistics*, 9:432–441.
- [59] Frisch, S. M., Schaller, M., and Cieply, B. (2013). Mechanisms that link the oncogenic epithelial–mesenchymal transition to suppression of anoikis. *Journal of Cell Science*, 126(1):21–29.
- [60] Gardiner-Garden, M. and Frommer, M. (1987). CpG islands in vertebrate genomes. *Journal of Molecular Biology*, 196(2):261–282.
- [61] Garofalo, M., Quintavalle, C., Romano, G., Croce, C., and Condorelli, G. (2012). miR221/222 in cancer: their role in tumor progression and response to therapy. *Current Molecular Medicine*, 12(1):27–33.
- [62] Geiger, T. and Levitzki, A. (2007). Loss of robustness and addiction to IGF1 during early keratinocyte transformation by human papilloma virus 16. *PLoS One*, 2:605.

- [63] Genovesi, L. A., Carter, K. W., Gottardo, N. G., Giles, K. M., and Dallas, P. B. (2011). Integrated analysis of miRNA and mRNA expression in childhood medulloblastoma compared with neural stem cells. *PloS One*, 6(9):e23935.
- [64] Guadamillas, M. C., Cerezo, A., and del Pozo, M. A. (2011). Overcoming anoikis pathways to anchorage-independent growth in cancer. *Journal of Cell Science*, 124:3189–3197.
- [65] Hamilton, J. D. (1994). *Time series analysis*. Princeton University Press.
- [66] Hanash, S. (2004). Integrated global profile of cancer. *Nature Reviews Cancer*, 4:638–644.
- [67] Harden, S. V., Guo, Z., Epstein, J. I., and Sidransky, D. (2003). Quantitative GSTP1 methylation clearly distinguishes benign prostatic tissue and limited prostate adenocarcinoma. *The Journal of Urology*, 169(3):1138–1142.
- [68] Harville, D. A. (2008). *Matrix Algebra From a Statistician's Perspective*. Springer, New York.
- [69] Heinen, A., Rengifo, E., et al. (2003). *Multivariate modelling of time series count data: an autoregressive conditional Poisson model*. Universite Catholique de Louvain.
- [70] Henken, F. E., Arce, J. D., Rösl, F., Bosch, L., Meijer, C. J. L. M., Snijders, P. J. F., and Steenbergen, R. D. M. (2012). The functional role of notch signaling in HPV-mediated transformation is dose-dependent and linked to AP-1 alterations. *Cellular Oncology*, 35(2):77–84.
- [71] Henken, F. E., Wilting, S. M., Overmeer, R. M., Van Rietschoten, J. G., Nygren, A. O. H., Errami, A., Schouten, J. P., Meijer, C. J. L. M., Snijders, P. J. F., and Steenbergen, R. D. M. (2007). Sequential gene promoter methylation during HPV-induced cervical carcinogenesis. *British Journal of Cancer*, 97:1457–1464.
- [72] Hong, F. and Li, H. (2006). Functional Hierarchical Models for Identifying Genes with Different Time-Course Expression Profiles. *Biometrics*, 62:534–544.
- [73] Hosseini, E. S., Kashani, H. H., Sabzalipoor, H., Nikzad, H., Meryet-Figuere, M., and Asemi, Z. (2017). Dysregulated expression of long noncoding RNAs in gynecologic cancers. *Molecular Cancer*, 16(1):107.
- [74] Hou, F., Li, Z., Ma, D., Zhang, W., Zhang, Y., Zhang, T., Kong, B., and Cui, B. (2012). Distribution of Th17 cells and Foxp3-expressing T cells in tumor-infiltrating lymphocytes in patients with uterine cervical cancer. *Clinica Chimica Acta*, 413(23):1848–1854.
- [75] Huang, Y. W., Liu, J. C., Deatherage, D. E., Luo, J., Mutch, D. G., Goodfellow, P. J., Miller, D. S., and Huang, T. H. M. (2009). Epigenetic repression of microRNA-129-2 leads to overexpression of SOX4 oncogene in endometrial cancer. *Cancer Research*, 69(23):9038–9046.
- [76] Huber, W., Von Heydebreck, A., Sültmann, H., Poustka, A., and Vingron, M. (2002). Variance stabilization applied to microarray data calibration and to the quantification of differential expression. *Bioinformatics*, 18(suppl 1):S96–S104.

- [77] Irizarry, R. A., Hobbs, B., Collin, F., Beazer-Barclay, Y. D., Antonellis, K. J., Scherf, U., Speed, T. P., et al. (2003). Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics*, 4(2):249–264.
- [78] Jacobs, M. V., Snijders, P. J., Van Den Brule, A. J., Helmerhorst, T. J., Meijer, C. J., and Walboomers, J. M. (1997). A general primer GP5+/GP6 (+)-mediated PCR-enzyme immunoassay method for rapid detection of 14 high-risk and 6 low-risk human papillomavirus genotypes in cervical scrapings. *Journal of Clinical Microbiology*, 35(3):791–795.
- [79] Jones, B. and West, M. (2005). Covariance decomposition in undirected Gaussian graphical models. *Biometrika*, 92(4):779–786.
- [80] Kamio, M., Yoshida, T., Ogata, H., Douchi, T., Nagata, Y., Inoue, M., Hasegawa, M., Yonemitsu, Y., and Yoshimura, A. (2004). SOX1 inhibits HPV-E7-mediated transformation by inducing degradation of E7 protein. *Oncogene*, 23:3107–3115.
- [81] Kanehisa, M. and Goto, S. (2000). KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Research*, 28(1):27–30.
- [82] Kantz, H. and Schreiber, T. (2004). *Nonlinear time series analysis*, volume 7. Cambridge University Press.
- [83] Khalifa, Y. B., Teissier, S., Tan, M. K. M., Phan, Q. T., Daynac, M., Wong, W. Q., and Thierry, F. (2011). The human papillomavirus E6 oncogene represses a cell adhesion pathway and disrupts focal adhesion through degradation of TAp63 β upon transformation. *PLoS Pathology*, 7:e1002256.
- [84] Klingelhutz, A. J. and Roman, A. (2012). Cellular transformation by human papillomaviruses: lessons learned by comparing high-and low-risk viruses. *Virology*, 424(2):77–98.
- [85] Kodama, J., Hashimoto, I., Seki, N., Hongo, A., Yoshinouchi, M., Okuda, H., and Kudo, T. (2001). Thrombospondin-1 and-2 messenger RNA expression in invasive cervical cancer correlation with angiogenesis and prognosis. *Clinical Cancer Research*, 7:2826–2831.
- [86] Korzeniewski, N., Spardy, N., Duensing, A., and Duensing, S. (2011). Genomic instability and cancer: lessons learned from human papillomaviruses. *Cancer Letters*, 305(2):113–122.
- [87] Kozaki, K., Imoto, I., Mogi, S., Omura, K., and Inazawa, J. (2008). Exploration of tumor-suppressive microRNAs silenced by DNA hypermethylation in oral cancer. *Cancer Research*, 68(7):2094–2105.
- [88] Kristensen, L. S. and Hansen, L. L. (2009). PCR-based methods for detecting single-locus DNA methylation biomarkers in cancer diagnostics, prognostics, and response to treatment. *Clinical Chemistry*, 55(8):1471–1483.
- [89] Langevin, S. M., Stone, R. A., Bunker, C. H., Grandis, J. R., Sobol, R. W., and Taioli, E. (2010). MicroRNA-137 promoter methylation in oral rinses from patients with squamous cell carcinoma of the head and neck is associated with gender and body mass index. *Carcinogenesis*, 31(5):864–870.

- [90] Langevin, S. M., Stone, R. A., Bunker, C. H., Lyons-Weiler, M. A., LaFramboise, W. A., Kelly, L., Seethala, R. R., Grandis, J. R., Sobol, R. W., and Taioli, E. (2011). MicroRNA-137 promoter methylation is associated with poorer overall survival in patients with squamous cell carcinoma of the head and neck. *Cancer*, 117(7):1454–1462.
- [91] Lauritzen, S. L. (1996). *Graphical models*. Oxford University Press, Oxford.
- [92] Lebre, S., Becq, J., Devaux, F., Stumpf, M. P. H., and Lelandais, G. (2010). Statistical inference of the time-varying structure of gene-regulation networks. *BMC Systems Biology*, 4(1):130.
- [93] Lindenbergh-Van der Plas, M., Martens-de Kemp, S. R., de Maaker, M., Van Wieringen, W. N., Ylstra, B., Agami, R., Cerisoli, F., Leemans, C. R., Braakhuis, B. J. M., and Brakenhoff, R. H. (2013). Identification of lethal microRNAs specific for head and neck cancer. *Clinical Cancer Research*, 19:5647–5657.
- [94] Liu, H., Han, F., Yuan, M., Lafferty, J., Wasserman, L., et al. (2012). High-dimensional semiparametric Gaussian copula graphical models. *The Annals of Statistics*, 40(4):2293–2326.
- [95] Liu, H., Lafferty, J., and Wasserman, L. (2009). The nonparanormal: Semiparametric estimation of high dimensional undirected graphs. *Journal of Machine Learning Research*, 10(Oct):2295–2328.
- [96] Livak, K. J. and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\delta\delta$ CT method. *Methods*, 25(4):402–408.
- [97] Long, X. H., Zhou, Y. F., Peng, A. F., Zhang, Z. H., Chen, X. Y., Chen, W. Z., Liu, J. M., Huang, S. H., and Liu, Z. L. (2015). Demethylation-mediated miR-129-5p up-regulation inhibits malignant phenotype of osteogenic osteosarcoma by targeting homo sapiens valosin-containing protein (VCP). *Tumor Biology*, 36(5):3799–3806.
- [98] Lönnstedt, I. and Speed, T. P. (2002). Replicated microarray data. *Statistica Sinica*, 12:31–46.
- [99] López, J. A. and Alvarez-Salas, L. M. (2011). Differential effects of miR-34c-3p and miR-34c-5p on SiHa cells proliferation apoptosis, migration and invasion. *Biochemical and Biophysical Research Communications*, 409:513–519.
- [100] Lu, C. Y., Lin, K. Y., Tien, M. T., Wu, C. T., Uen, Y. H., and Tseng, T. L. (2013). Frequent DNA methylation of miR-129-2 and its potential clinical implication in hepatocellular carcinoma. *Genes, Chromosomes and Cancer*, 52(7):636–643.
- [101] Lujambio, A., Calin, G. A., Villanueva, A., Ropero, S., Sánchez-Céspedes, M., Blanco, D., Montuenga, L. M., Rossi, S., Nicoloso, M. S., Faller, W. J., et al. (2008). A microRNA DNA methylation signature for human cancer metastasis. *Proceedings of the National Academy of Sciences*, 105(36):13556–13561.
- [102] Luo, D., Wang, J., Li, J., and Post, M. (2011). Mouse snail is a target gene for HIF. *Molecular Cancer Research*, 9(2):234–245.

- [103] Lütkepohl, H. (2005). *New introduction to multiple time series analysis*. Springer.
- [104] Mack, H. I. D. and Munger, K. (2013). The LKB1 tumor suppressor differentially affects anchorage independent growth of HPV positive cervical cancer cell lines. *Virology*, 446:9–16.
- [105] Massagué, J. (2008). TGF- β in cancer. *Cell*, 134(2):215–230.
- [106] McCormack, S. J., Brazinski, S. E., Moore Jr, J. L., Werness, B. A., and Goldstein, D. J. (1997). Activation of the focal adhesion kinase signal transduction pathway in cervical carcinoma cell lines and human genital epithelial cells immortalized with human papillomavirus type 18. *Oncogene*, 15(3).
- [107] Medina-Martinez, I., Barrón, V., Roman-Bassaure, E., Juárez-Torres, E., Guardado-Estrada, M., Espinosa, A., Bermudez, M., Fernández, F., Venegas-Vega, C., Orozco, L., et al. (2014). Impact of gene dosage on gene expression, biological processes and survival in cervical cancer: a genome-wide follow-up study. *PLoS One*, 9(5):e97842.
- [108] Mersmann, O. (2014). microbenchmark: Accurate timing functions. *CRAN*, 1.4-2:12.
- [109] Meyer, P. E., Kontos, K., Lafitte, F., and Bontempi, G. (2007). Information-theoretic inference of large transcriptional regulatory networks. *EURASIP Journal on Bioinformatics and Systems Biology*, 2007(1):79879.
- [110] Miok, V. (2018). Supplementary materials: Comprehensive molecular characterization of HPV-induced transformation by longitudinal statistical modelling. <https://github.com/viktormiok/PhdThesisSupplementaryMaterials>, 1:1–96.
- [111] Miok, V., Wilting, S. M., van de Wiel, M. A., Jaspers, A., van Noort, P. I., Brakenhoff, R. H., Snijders, P. J. F., Steenbergen, R., and van Wieringen, W. N. (2014). tigar: integrative significance analysis of temporal differential gene expression induced by genomic abnormalities. *BMC Bioinformatics*, 15(1):327.
- [112] Miok, V., Wilting, S. M., and van Wieringen, W. N. (2017). Ridge estimation of the VAR(1) model and its time series chain graph from multivariate time-course omics data. *Biometrical Journal*, 59(1):172–191.
- [113] Miranda, K. C., Huynh, T., Tay, Y., Ang, Y. S., Tam, W. L., Thomson, A. M., Lim, B., and Rigoutsos, I. (2006). A pattern-based method for the identification of microRNA binding sites and their corresponding heteroduplexes. *Cell*, 126(6):1203–1217.
- [114] Monti, S., Tamayo, P., Mesirov, J., and Golub, T. (2003). Consensus clustering: a resampling-based method for class discovery and visualization of gene expression microarray data. *Machine Learning*, 52(1-2):91–118.
- [115] Moody, C. A. and Laimins, L. A. (2010). Human papillomavirus oncoproteins: pathways to transformation. *Nature Reviews Cancer*, 10:550–560.

- [116] Mori, S., Chang, J. T., Andrechek, E. R., Matsumura, N., Baba, T., Yao, G., Kim, J. W., Gatza, M., Murphy, S., and Nevins, J. R. (2009). Anchorage-independent cell growth signature identifies tumors with metastatic potential. *Oncogene*, 28:2796–2805.
- [117] Mutarelli, M., Cicatiello, L., Ferraro, L., Grober, O. M. V., Ravo, M., Facchiano, A. M., Angelini, C., and Weisz, A. (2008). Time-course analysis of genome-wide gene expression data from hormone-responsive human breast cancer cells. *BMC Bioinformatics*, 9:S12.
- [118] Nau, G. J., Richmond, J. F. L., Schlesinger, A., Jennings, E. G., Lander, E. S., and Young, R. A. (2002). Human macrophage activation programs induced by bacterial pathogens. *Proceedings of the National Academy of Sciences*, 99:1503–1508.
- [119] Neuzillet, C., Tijeras-Raballand, A., Cohen, R., Cros, J., Faivre, S., Raymond, E., and de Gramont, A. (2015). Targeting the TGF β pathway for cancer therapy. *Pharmacology & Therapeutics*, 147:22–31.
- [120] Newman, M. (2010). *Networks: an introduction*. Oxford University Press.
- [121] Nicholls, D. F. and Pope, A. L. (1988). Bias in the estimation of multivariate autoregressions. *Australian Journal of Statistics*, 30A:296–309.
- [122] Novianti, P. W., Snoek, B. C., Wilting, S. M., and van de Wiel, M. A. (2017). Better diagnostic signatures from RNAseq data through use of auxiliary co-data. *Bioinformatics*, 33(10):1572–1574.
- [123] Oberhofer, W. and Kmenta, J. (1974). A general procedure for obtaining maximum likelihood estimates in generalized regression models. *Econometrica*, 42:579–590.
- [124] Olshen, A. B., Venkatraman, E. S., Lucito, R., and Wigler, M. (2004). Circular binary segmentation for the analysis of array-based DNA copy number data. *Biostatistics*, 5(4):557–572.
- [125] Oshlack, A., Robinson, M. D., and Young, M. D. (2010). From RNA-seq reads to differential expression results. *Genome Biology*, 11:220.
- [126] Overmeer, R. M., Henken, F. E., Bierkens, M., Wilting, S. M., Timmerman, I., Meijer, C. J. L. M., Snijders, P. J. F., and Steenbergen, R. D. M. (2009). Repression of mal tumour suppressor activity by promoter methylation during cervical carcinogenesis. *The Journal of Pathology*, 219(3):327–336.
- [127] Overmeer, R. M., Henken, F. E., Snijders, P. J. F., Claassen-Kramer, D., Berkhof, J., Helmerhorst, T. J. M., Heideman, D. A. M., Wilting, S. M., Murakami, Y., Ito, A., Meijer, C. J. L. M., and Steenbergen, R. D. M. (2008). Association between dense CADM1 promoter methylation and reduced protein expression in high-grade CIN and cervical SCC. *The Journal of Pathology*, 215:388–397.
- [128] Paoli, P., Giannoni, E., and Chiarugi, P. (2013). Anoikis molecular pathways and its role in cancer progression. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research*, 1833(12):3481–3498.

- [129] Park, N. H., Min, B. M., Li, S. L., Huang, M. Z., and Doniger, J. (1991). Immortalization of normal human oral keratinocytes with type 16 human papillomavirus. *Carcinogenesis*, 12(9):1627–1631.
- [130] Pecoraro, G., Morgan, D., and Defendi, V. (1989). Differential effects of human papillomavirus type 6, 16, and 18 DNAs on immortalization and transformation of human cervical epithelial cells. *Proceedings of the National Academy of Sciences*, 86(2):563–567.
- [131] Peel, D. and McLachlan, G. J. (2000). Robust mixture modeling using the t distribution. *Statistics and Computing*, 10:339–348.
- [132] Pett, M. R., Alazawi, W. O. F., Roberts, I., Downen, S., Smith, D. I., Stanley, M. A., and Coleman, N. (2004). Acquisition of high-level chromosomal instability is associated with integration of human papillomavirus type 16 in cervical keratinocytes. *Cancer research*, 64(4):1359–1368.
- [133] Pinzón, N., Li, B., Martinez, L., Sergeeva, A., Presumey, J., Apparailly, F., and Seitz, H. (2017). microRNA target prediction programs predict many false positives. *Genome Research*, 27(2):234–245.
- [134] Pirisi, L., Yasumoto, S., Feller, M., Doniger, J., and DiPaolo, J. A. (1987). Transformation of human fibroblasts and keratinocytes with human papillomavirus type 16 DNA. *Journal of Virology*, 61(4):1061–1066.
- [135] Rau, A., Jaffrézic, F., Foulley, J. L., and Doerge, R. W. (2010). An empirical Bayesian method for estimating biological networks from temporal microarray data. *Statistical Applications in Genetics and Molecular Biology*, 9(1).
- [136] Razani, B., Altschuler, Y., Zhu, L., Pestell, R. G., Mostov, K. E., and Lisanti, M. P. (2000). Caveolin-1 expression is down-regulated in cells transformed by the human papilloma virus in a p53-dependent manner. Replacement of caveolin-1 expression suppresses HPV-mediated cell transformation. *Biochemistry*, 39:13916–13924.
- [137] Ritchie, M. E., Silver, J., Oshlack, A., Holmes, M., Diyagama, D., Holloway, A., and Smyth, G. K. (2007). A comparison of background correction methods for two-colour microarrays. *Bioinformatics*, 23:2700–2707.
- [138] Robinson, M. D., McCarthy, D. J., and Smyth, G. K. (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*, 26:139–140.
- [139] Robinson, M. D. and Smyth, G. K. (2007). Moderated statistical tests for assessing differences in tag abundance. *Bioinformatics*, 23:2881–2887.
- [140] Rue, H., Matrino, S., and Chopin, N. (2009). Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximation. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 71:319–392.
- [141] Ruiz-Lafuente, N., Alcaraz-García, M. J., García-Serna, A. M., Sebastián-Ruiz, S., Moya-Quiles, M. R., García-Alonso, A. M., and Parrado, A. (2015). Dock10, a Cdc42 and Rac1 GEF, induces loss of elongation, filopodia, and ruffles in cervical cancer epithelial HeLa cells. *Biology Open*, 4(5):627–635.

- [142] Ruppert, D., Wand, M. P., and Carroll, R. J. (2009). Semiparametric regression during 2003-2007. *Electronic Journal of Statistics*, 3:1193.
- [143] Sand, M., Skrygan, M., Sand, D., Georgas, D., Gambichler, T., Hahn, S. A., Altmeyer, P., and Bechara, F. G. (2013). Comparative microarray analysis of microRNA expression profiles in primary cutaneous malignant melanoma, cutaneous malignant melanoma metastases, and benign melanocytic nevi. *Cell and Tissue Research*, 351(1):85–98.
- [144] Schena, M., Shalon, D., Davis, R. W., and Brown, P. O. (1995). Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science*, 270:467.
- [145] Schmittgen, T. D. and Livak, K. J. (2008). Analyzing real-time PCR data by the comparative CT method. *Nature Protocols*, 3(6):1101–1108.
- [146] Schütze, D. M., Krijgsman, O., Snijders, P. J. F., Ylstra, B., Weischenfeldt, J., Mardin, B., Stütz, A. M., Korb, J. O., de Winter, J. P., Meijer, C. J. L. M., et al. (2016). Immortalization capacity of HPV types is inversely related to chromosomal instability. *Oncotarget*, 7(25):37608.
- [147] Schütze, D. M., Snijders, P. J. F., Bosch, L., Kramer, D., Meijer, C. J. L. M., and Steenbergen, R. D. M. (2014). Differential in vitro immortalization capacity of eleven, probable high-risk human papillomavirus types. *Journal of Virology*, 88(3):1714–1724.
- [148] Schwock, J., Dhani, N., Cao, M. P. J., Zheng, J., Clarkson, R., Radulovich, N., Navab, R., Horn, L.-C., and Hedley, D. W. (2009). Targeting focal adhesion kinase with dominant-negative FRNK or Hsp90 inhibitor 17-DMAG suppresses tumor growth and metastasis of SiHa cervical xenografts. *Cancer Research*, 69(11):4750–4759.
- [149] Shapira, S. D., Gat-Viks, I., Shum, B. O. V., Dricot, A., de Grace, M. M., Wu, L., Gupta, P. B., Hao, T., Silver, S. J., Root, D. E., et al. (2009). A physical and regulatory map of host-influenza interactions reveals pathways in H1N1 infection. *Cell*, 139(7):1255–1267.
- [150] Shen, R., Pan, S., Qi, S., Lin, X., and Cheng, S. (2010). Epigenetic repression of microRNA-129-2 leads to overexpression of SOX4 in gastric cancer. *Biochemical and Biophysical Research Communications*, 394(4):1047–1052.
- [151] Shen, W., Tao, G. G., Zhang, Y., Cai, B., Sun, J., and Tian, Z. G. (2017). TGF- β in pancreatic cancer initiation and progression: two sides of the same coin. *Cell & Bioscience*, 7(1):39.
- [152] Shimizu, T., Suzuki, H., Nojima, M., Kitamura, H., Yamamoto, E., Maruyama, R., Ashida, M., Hatahira, T., Kai, M., Masumori, N., et al. (2013). Methylation of a panel of microRNA genes is a novel biomarker for detection of bladder cancer. *European Urology*, 63(6):1091–1100.
- [153] Silber, J., Lim, D. A., Petritsch, C., Persson, A. I., Maunakea, A. K., Yu, M., Vandenberg, S. R., Ginzinger, D. G., James, C. D., Costello, J. F., et al. (2008).

- miR-124 and miR-137 inhibit proliferation of glioblastoma multiforme cells and induce differentiation of brain tumor stem cells. *BMC Medicine*, 6(1):1.
- [154] Silver, J. D., Ritchie, M. E., and Smyth, G. K. (2008). Microarray background correction: maximum likelihood estimation for the normal exponential convolution. *Biostatistics*, 10(2):352–363.
- [155] Smeets, S. J., van der Plas, M., Schaaij-Visser, T., van Veen, E. A. M., van Meerloo, J., Braakhuis, B. J. M., Steenbergen, R. D. M., and Brakenhoff, R. H. (2011). Immortalization of oral keratinocytes by functional inactivation of the p53 and pRb pathways. *International Journal of Cancer*, 128(7):1596–1605.
- [156] Smyth, G. K. (2004). Linear models and empirical Bayes methods for assessing differential expression in microarray experiments. *Statistical Applications in Genetics and Molecular Biology*, 3(1):1–25.
- [157] Snellenberg, S., Cillessen, S. A. G. M., Van Criekinge, W., Bosch, L., Meijer, C. J. L. M., Snijders, P. J. F., and Steenbergen, R. D. M. (2014). Methylation-mediated repression of PRDM14 contributes to apoptosis evasion in HPV-positive cancers. *Carcinogenesis*, 35(11):2611–2618.
- [158] Snellenberg, S., De Strooper, L. M. A., Hesselink, A. T., Meijer, C. J. L. M., Snijders, P. J. F., Heideman, D. A. M., and Steenbergen, R. D. M. (2012). Development of a multiplex methylation-specific PCR as candidate triage test for women with an HPV-positive cervical scrape. *BMC Cancer*, 12(1):1.
- [159] Song, L., Kolar, M., and Xing, E. P. (2009). Time-varying dynamic Bayesian networks. pages 1732–1740. Elsevier.
- [160] Sopov, I., Sørensen, T., Magbagbeolu, M., Jansen, L., Beer, K., Kühne-Heid, R., Kirchmayr, R., Schneider, A., and Dürst, M. (2004). Detection of cancer-related gene expression profiles in severe cervical neoplasia. *International Journal of Cancer*, 112(1):33–43.
- [161] Soto-Reyes, E., González-Barrios, R., Cisneros-Soberanis, F., Herrera-Goepfert, R., Pérez, V., Cantú, D., Prada, D., Castro, C., Recillas-Targa, F., and Herrera, L. A. (2012). Disruption of CTCF at the miR-125b1 locus in gynecological cancers. *BMC Cancer*, 12(1):1.
- [162] Spiegelhalter, D. J., Best, N. G., Carlin, B. P., and van der Liner, A. (2002). Bayesian measures of model complexity and fit. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 64:583–639.
- [163] Srivastava, K., Pickard, A., McDade, S., and McCance, D. J. (2015). p63 drives invasion in keratinocytes expressing HPV16 E6/E7 genes through regulation of Src-FAK signalling. *Oncotarget*.
- [164] Srivastava, S. K., Ahmad, A., Zubair, H., Miree, O., Singh, S., Rocconi, R. P., Scalici, J., and Singh, A. P. (2017). MicroRNAs in gynecological cancers: Small molecules with big implications. *Cancer Letters*.

- [165] Steenbergen, R. D. M., de Wilde, J., Wilting, S. M., Brink, A. A. T. P., Snijders, P. J. F., and Meijer, C. J. L. M. (2005). HPV-mediated transformation of the anogenital tract. *Journal of Clinical Virology*, 32:25–33.
- [166] Steenbergen, R. D. M., Kramer, D., Braakhuis, B. J. M., Stern, P. L., Verheijen, R. H. M., Meijer, C. J. L. M., and Snijders, P. J. F. (2004). TSLC1 gene silencing in cervical cancer cell lines and cervical neoplasia. *Journal of the National Cancer Institute*, 96:294–305.
- [167] Steenbergen, R. D. M., Kramer, D., Meijer, C. J. L. M., Walboomers, J. M. M., Trott, D. A., Cuthbert, A. P., Newbold, R. F., Overkamp, W. J. I., Zdzienicka, M. Z., and Snijders, P. J. F. (2001). Telomerase suppression by chromosome 6 in a human papillomavirus type 16-immortalized keratinocyte cell line and in a cervical cancer cell line. *Journal of the National Cancer Institute*, 93(11):865–872.
- [168] Steenbergen, R. D. M., Snijders, P. J. F., Heideman, D. A. M., and Meijer, C. J. L. M. (2014). Clinical implications of (epi) genetic changes in HPV-induced cervical precancerous lesions. *Nature Reviews Cancer*, 14:395–405.
- [169] Steenbergen, R. D. M., Walboomers, J. M. M., Meijer, C. J. L. M., Van der Raaij-Helmer, E. M. H., Parker, J. N., Chow, L. T., Broker, T. R., and Snijders, P. J. F. (1996). Transition of human papillomavirus type 16 and 18 transfected human foreskin keratinocytes towards immortality: activation of telomerase and allele loss at 3p, 10p, 11q and/or 18q. *Oncogene*, 13:1249–1257.
- [170] Steponaitiene, R., Kupcinskas, J., Langner, C., Balaguer, F., Venclauskas, L., Pauzas, H., Tamelis, A., Skieceviciene, J., Kupcinskas, L., Malfertheiner, P., et al. (2015). Epigenetic silencing of miR-137 is a frequent event in gastric carcinogenesis. *Molecular Carcinogenesis*.
- [171] Storey, J. D., Xiao, W., Leek, J. T., Tompkins, R. G., and Davis, R. W. (2005). Significance analysis of time course microarray experiments. *Proceedings of the National Academy of Sciences of the United States of America*, 102:12837–12842.
- [172] Strimmer, K. (2008). A unified approach to false discovery rate estimation. *BMC Bioinformatics*, 9(1):303.
- [173] Sun, L., Li, T., Wei, Q., Zhang, Y., Jia, X., Wan, Z., and Han, L. (2014). Upregulation of BNIP3 mediated by ERK/HIF-1 α pathway induces autophagy and contributes to anoikis resistance of hepatocellular carcinoma cells. *Future Oncology*, 10(8):1387–1398.
- [174] Szalmás, A. and Kónya, J. (2009). Epigenetic alterations in cervical carcinogenesis. 19:144–152.
- [175] Tabas-Madrid, D., Muniategui, A., Sánchez-Caballero, I., Martínez-Herrera, D. J., Sorzano, C. O. S., Rubio, A., and Pascual-Montano, A. (2014). Improving miRNA-mRNA interaction predictions. *BMC Genomics*, 15(10):S2.
- [176] Tai, Y. C. and Speed, T. P. (2006). A multivariate empirical Bayes statistics for replicated microarray time course data. *The Annals of Statistics*, 34:2387–2412.

- [177] Takwi, A. A., Wang, Y. M., Wu, J., Michaelis, M., Cinatl, J., and Chen, T. (2014). miR-137 regulates the constitutive androstane receptor and modulates doxorubicin sensitivity in parental and doxorubicin-resistant neuroblastoma cells. *Oncogene*, 33(28):3717–3729.
- [178] Tang, W., Feng, X., Zhang, S., Ren, Z., Liu, Y., Yang, B., Cai, Y., Xia, J., Ge, N., et al. (2015). Caveolin-1 confers resistance of hepatoma cells to anoikis by activating IGF-1 pathway. *Cellular Physiology and Biochemistry*, 36(3):1223–1236.
- [179] Thomas, L. K., Bermejo, J. L., Vinokurova, S., Jensen, K., Bierkens, M., Steenberg, R., Bergmann, M., von Knebel Doeberitz, M., and Reuschenbach, M. (2014). Chromosomal gains and losses in human papillomavirus-associated neoplasia of the lower genital tract—a systematic review and meta-analysis. *European Journal of Cancer*, 50(1):85–98.
- [180] Touleimat, N. and Tost, J. (2012). Complete pipeline for Infinium® Human Methylation 450K BeadChip data processing using subset quantile normalization for accurate DNA methylation estimation. *Epigenomics*, 4(3):325–341.
- [181] Toussaint-Smith, E. D. D. B. and Roman, A. (2004). Expression of human papillomavirus type 16 E6 and E7 oncoproteins in primary foreskin keratinocytes is sufficient to alter the expression of angiogenic factors. *Oncogene*, 23:2988–2995.
- [182] Tsai, K. W., Wu, C. W., Hu, L. Y., Li, S. C., Liao, Y. L., Lai, C. H., Kao, H. W., Fang, W. L., Huang, K. H., Chan, W. C., et al. (2011). Epigenetic regulation of miR-34b and miR-129 expression in gastric cancer. *International Journal of Cancer*, 129(11):2600–2610.
- [183] Tusher, V. G., Tibshirani, R., and Chu, G. (2001). Significance analysis of microarrays applied to the ionizing radiation response. *Proceedings of the National Academy of Sciences*, 98(9):5116–5121.
- [184] Van de Wiel, M. A., Kim, K. I., Vosse, S. J., Van Wieringen, W. N., Wilting, S. M., and Ylstra, B. (2007). CGHcall: calling aberrations for array CGH tumor profiles. *Bioinformatics*, 23:892–894.
- [185] Van de Wiel, M. A., Leday, G. G. R., Pardo, L., Rue, H., Van der Vaart, A. W., and Van Wieringen, W. N. (2013). Bayesian analysis of RNA sequencing data by estimating multiple shrinkage priors. *Biostatistics*, 14:113–128.
- [186] Van de Wiel, M. A., Picard, F., Van Wieringen, W. N., and Ylstra, B. (2010). Preprocessing and downstream analysis of microarray DNA copy number profiles. *Briefings in Bioinformatics*, 12(1):10–21.
- [187] Van Loan, C. F. (2000). The ubiquitous kronecker product. *Journal of Computational and Applied Mathematics*, 123:85–100.
- [188] Van Wieringen, W. N., Berkhof, J., and Van de Wiel, M. A. (2010). A random coefficients model for regional co-expression associated with DNA copy number. *Statistical Applications in Genetics and Molecular Biology*, 9:1–30.

- [189] Van Wieringen, W. N. and Peeters, C. F. W. (2016). Ridge estimation of inverse covariance matrices from high-dimensional data. *Computational Statistics & Data Analysis*, 103:284–303.
- [190] Van Wieringen, W. N., Unger, K., Leday, G. G. R., Krijgsman, O., de Menezes, R., Ylstra, B., and Van de Wiel, M. A. (2012). Matching of array CGH and gene expression microarray features for the purpose of integrative genomic analysis. *BMC Bioinformatics*, 13:80.
- [191] Van Wieringen, W. N., Van de Wiel, M. A., and Ylstra, B. (2007). Weighted clustering of called array CGH data. *Biostatistics*, 9(3):484–500.
- [192] Van Wieringen, W. N. and van der Vaart, A. W. (2015). Transcriptomic heterogeneity in cancer as a consequence of dysregulation of the gene-gene interaction network. *Bulletin of Mathematical Biology*, 77(9):1768–1786.
- [193] Van Zeeburg, H. J. T., Snijders, P. J. F., Pals, G., Hermsen, M. A., Rooimans, M. A., Bagby, G., Soulier, J., Gluckman, E., Wennerberg, J., Leemans, C. R., et al. (2005). Generation and molecular characterization of head and neck squamous cell lines of fanconi anemia patients. *Cancer Research*, 65(4):1271–1276.
- [194] Veigaard, C. and Kjeldsen, E. (2014). Exploring the genome-wide relation between copy number status and microRNA expression. *Genomics*, 104(4):271–278.
- [195] Vogelstein, B. and Kinzler, K. W. (2004). Cancer genes and the pathways they control. *Nature Medicine*, 10:789–799.
- [196] Vrba, L., Muñoz-Rodríguez, J. L., Stampfer, M. R., and Futscher, B. W. (2013). miRNA gene promoters are frequent targets of aberrant DNA methylation in human breast cancer. *PloS One*, 8(1):e54398.
- [197] Walboomers, J. M. M., Jacobs, M. V., Manos, M. M., Bosch, F. X., Kummer, J. A., Shah, K. V., Snijders, P. J. F., Peto, J., Meijer, C. J. L. M., and Muñoz, N. (1999). Human papillomavirus is a necessary cause of invasive cervical cancer worldwide. *The Journal of Pathology*, 189(1):12–19.
- [198] Wang, F., Liu, M., Li, X., and Tang, H. (2013). MiR-214 reduces cell survival and enhances cisplatin-induced cytotoxicity via down-regulation of Bcl2l2 in cervical cancer cells. *FEBS Letters*, 587(5):488–495.
- [199] Wang, Z., Gerstein, M., and Snyder, M. (2009). RNA-Seq: a revolutionary tool for transcriptomics. *Nature Reviews Genetics*, 10:57–63.
- [200] Weinberg, R. A. (2006). *The Biology of Cancer*. Garland Science, New York.
- [201] Wentzensen, N., Sherman, M. E., Schiffman, M., and Wang, S. S. (2009). Utility of methylation markers in cervical cancer early detection: appraisal of the state-of-the-science. *Gynecologic Oncology*, 112:293–299.
- [202] Whittaker, J. (1990). *Graphical Models in Applied Multivariate Statistics*. John Wiley & Sons, Chichester, England.

- [203] Wiklund, E. D., Gao, S., Hulf, T., Sibbritt, T., Nair, S., Costea, D. E., Villadsen, S. B., Bakholdt, V., Bramsen, J. B., Sørensen, J. A., et al. (2011). MicroRNA alterations and associated aberrant DNA methylation patterns across multiple sample types in oral squamous cell carcinoma. *PLoS One*, 6(11):e27840.
- [204] Willey, J. C., Broussoud, A., Sleemi, A., Bennett, W. P., Cerutti, P., and Harris, C. C. (1991). Immortalization of normal human bronchial epithelial cells by human papillomaviruses 16 or 18. *Cancer Research*, 51(19):5370–5377.
- [205] Wilting, S. M., de Wilde, J., Meijer, C. J. L., Berkhof, J., Yi, Y., Van Wieringen, W. N., Braakhuis, B. J. M., Meijer, G. A., Ylstra, B., Snijders, P. J. F., and Steenbergen, R. D. M. (2008). Integrated genomic and transcriptional profiling identifies chromosomal loci with altered gene expression in cervical cancer. *Genes, Chromosomes and Cancer*, 47:890–905.
- [206] Wilting, S. M., Snijders, P. J. F., Meijer, G. A., Ylstra, B., Van den Ijssel, P. R., Snijders, A. M., Albertson, D. G., Coffa, J., Schouten, J. P., Van de Wiel, M. A., Meijer, C. J. L. M., and Steenbergen, R. D. M. (2006). Increased gene copy number at chromosome 20q are frequent in both squamous cell carcinomas and adenocarcinomas of the cervix. *The Journal of Pathology*, 209:220–230.
- [207] Wilting, S. M. and Steenbergen, R. D. M. (2016). Molecular events leading to HPV-induced high grade neoplasia. *Papillomavirus Research*, 2:85–88.
- [208] Wilting, S. M., van Boerdonk, R. A. A., Henken, F. E., Meijer, C. J. L. M., Diosdado, B., Meijer, G. A., le Sage, C., Agami, R., Snijders, P. J. F., and Steenbergen, R. D. M. (2010). Methylation-mediated silencing and tumour suppressive function of hsa-miR-124 in cervical cancer. *Molecular Cancer*, 9(1):1.
- [209] Wilting, S. M., Verlaat, W., Jaspers, A., Makazaji, N. A., Agami, R., Meijer, C. J. L. M., Snijders, P. J. F., and Steenbergen, R. D. M. (2013). Methylation-mediated transcriptional repression of microRNAs during cervical carcinogenesis. *Epigenetics*, 8:220–228.
- [210] Wong, N. and Wang, X. (2014). miRDB: an online resource for microRNA target prediction and functional annotations. *Nucleic Acids Research*, 43(D1):D146–D152.
- [211] Wong, Y. F., Cheung, T. H., Tsao, G. S. W., Lo, K. W. K., Yim, S. F., Wang, V. W., Heung, M., Chan, S., Chan, L. K. Y., Ho, T. W. F., et al. (2006). Genome-wide gene expression profiling of cervical cancer in Hong Kong women by oligonucleotide microarray. *International Journal of Cancer*, 118(10):2461–2469.
- [212] Wu, H. S., Li, Y. F., Chou, C. I., Yuan, C. C., Hung, M. W., and Tsai, L. C. (2002). The concentration of serum Transforming Growth Factor beta-1 (TGF- β 1) is decreased in cervical carcinoma patients. *Cancer Investigation*, 20(1):55–59.
- [213] Yang, M. H., Wu, M. Z., Chiou, S. H., Chen, P. M., Chang, S. Y., Liu, C. J., Teng, S. C., and Wu, K. J. (2008). Direct regulation of TWIST by HIF-1 α promotes metastasis. *Nature Cell Biology*, 10(3):295–305.
- [214] Yao, T., Rao, Q., Liu, L., Zheng, C., Xie, Q., Liang, J., and Lin, Z. (2013). Exploration of tumor-suppressive microRNAs silenced by DNA hypermethylation in cervical cancer. *Virology Journal*, 10(1):1.

- [215] Ye, H., Zhang, Y., Geng, L., and Li, Z. (2015). Cdc42 expression in cervical cancer and its effects on cervical tumor invasion and migration. *International Journal of Oncology*, 46(2):757–763.
- [216] Zellner, A. (1962). An efficient method of estimating seemingly unrelated regressions and tests for aggregation bias. *Journal of the American Statistical Association*, 57:977–992.
- [217] Zhang, J., Li, S., Yan, Q., Chen, X., Yang, Y., Liu, X., and Wan, X. (2013). Interferon- β induced microRNA-129-5p down-regulates HPV-18 E6 and E7 viral gene expression by targeting SP1 in cervical cancer cells. *PLoS One*, 8(12):e81366.
- [218] Zhang, L., Huang, J., Yang, N., Greshock, J., Megraw, M., Giannakakis, A., Liang, S., Naylor, T., Barchetti, A., Ward, M., et al. (2006). microRNAs exhibit high frequency genomic alterations in human cancer. *Proceedings of the National Academy of Sciences*, 103(24):9136–9141.
- [219] Zhang, L., Volinia, S., Bonome, T., Calin, G. A., Greshock, J., Yang, N., Liu, C. G., Giannakakis, A., Alexiou, P., Hasegawa, K., et al. (2008). Genomic and epigenetic alterations deregulate microRNA expression in human epithelial ovarian cancer. *Proceedings of the National Academy of Sciences*, 105(19):7004–7009.
- [220] Zhu, H., Luo, H., Shen, Z., Hu, X., Sun, L., and Zhu, X. (2016). Transforming growth factor- β 1 in carcinogenesis, progression, and therapy in cervical cancer. *Tumor Biology*, 37(6):7075–7083.
- [221] Zhu, X., Er, K., Mao, C., Yan, Q., Xu, H., Zhang, Y., Zhu, J., Cui, F., Zhao, W., and Shi, H. (2013). miR-203 suppresses tumor growth and angiogenesis by targeting VEGFA in cervical cancer. *Cellular Physiology and Biochemistry*, 32(1):64–73.
- [222] Zoppoli, P., Morganella, S., and Ceccarelli, M. (2010). Timedelay-ARACNE: Reverse engineering of gene networks from time-course data by an information theoretic approach. *BMC Bioinformatics*, 11(1):154.

Summary

The body of all humans is built of small building blocks called cells, that carry out the functions necessary for life. The genetic material is contained within the nucleus of the cells. In the nucleus of the cell two copies of each of the 22 autosomal chromosomes and 2 sex chromosomes are stored. These chromosomes are composed of twisted ladder shaped deoxyribonucleic acid (DNA) containing 4 nucleotide bases called adenine (A), thymine (T), guanine (G), and cytosine (C). Genes are a specific combination of these nucleotides which together comprise a complete set of instructions for the cell to be able to function, grow and survive. In a multi-step process the information coded in the DNA is converted into a functional protein molecule that can be used by the cell. The first step in this process is transcription which uses information from the DNA to make messenger RNA (mRNA). In the second step, information from the mRNA is translated into a protein. In addition, several types of so-called non-coding RNA molecules are encoded by the DNA. These molecules are transcribed into RNA but not translated into protein. An example of these non-coding molecules are microRNAs (miRNAs). Instead of becoming protein, miRNAs bind to mRNAs and inhibit their translation into protein. In this thesis we investigated changes on DNA, mRNA and miRNA molecule levels.

Cancer is caused by a variety of factors that lead to abnormalities in DNA sequence and/or amount that change the normal functioning of the cell. Accumulation of these abnormalities and deregulations can ultimately lead to cancer. During this process of carcinogenesis, the genes that control the normal function of the cell (tumor suppressor genes) become inactive, while genes that initiate proliferation (oncogenes), start the process of fast and uncontrolled cell division. The process of carcinogenesis can be caused by various carcinogens like smoking, UV light, hereditary genetic alterations and viral infection. The most well-known case of viral infection caused carcinogenesis is cervical cancer, induced by human papillomavirus (HPV). Although more than 100 types of HPV exist, more than 70% of cervical carcinomas are caused by the high-risk types HPV16 and HPV18. Worldwide, cervical cancer is the fourth most diagnosed cancer in females, with the highest incidence in developing countries, due to the lack of population based screening programs. It is now widely accepted that involvement of high-risk types of HPV followed by genetic alterations can ultimately lead to cervical cancer.

The development and progression of cancer is a dynamic and complex biological process. To properly address the complexity of carcinogenesis, we need to simulate this process using experimental conditions we can control that allow us to follow this process over time. In this thesis we used a model consisting of normal human cells in which we introduced HPV that we interrogated at multiple moments in time at 3 molecular levels. In our cervical cancer experimental model two cell lines are infected with HPV16 and two with HPV18. The obtained cell line experimental model is shown to faithfully mimic cervical cancer development morphologically and

(epi)genetically, which is called transformation. During this transformation process we monitored all cell lines at different stages which allowed us to gain insights in genes consistently altered over time, as well as deregulation of a group of genes (pathways) that work together.

The goal of this thesis was to conduct comprehensive investigation of HPV-induced carcinogenesis using the above described model. For this we needed to develop and apply novel statistical methodology. The cell lines were profiled at 8 consecutive time points for DNA copy number, mRNA, and miRNA gene expression. Although many statistical models exist for analysis of time course data, all of them are focused on a single molecular level. However, we strongly believe that to be able to truly understand cancer, alterations in both DNA and RNA need to be investigated. So due to the fact that our experiment comprises data from different molecular levels, we needed to develop novel integrative statistical methodology to address this problem.

In this thesis we developed the integrative longitudinal statistical methodology for analyzing the data, both for single genes and groups of related genes (called pathways). In Chapter 2, which aims to perform temporal differential expression analysis, genes with the highest variation over time caused by abnormalities in DNA copy number were identified. On the other hand, analysis of well-defined groups of related genes (pathways) requires a methodology for network reconstruction presented in Chapter 3 and Chapter 4. Network reconstruction aims to identify gene-gene interactions between mRNAs, as well as relations between different molecular levels (e.g. how does a DNA copy number change affect this gene's mRNA expression levels). Employing this methodology to our experiment allows us to zoom-into the data and identify key genes within pathways that may be potential biomarkers or therapeutic targets.

Subsequent interpretation of the analysis result is divided into several steps, to ultimately reduce the number of potential gene candidates for an improved understanding of the underlying carcinogenic process. First, temporal differential mRNA and miRNA gene expression analysis were performed, where 106 miRNAs and 3642 mRNAs are identified with significant variation over time. Out of these number of genes, it is found that approximately 33% of differently expressed mRNA and miRNA are associated with chromosome abnormalities. All these genes are either up- or down-regulated in at least three out of four cell lines over time due to the presence of more or less DNA encoding for these genes. In order to further understand the effect of the observed changes for the functioning of the cell, the analysis is moved to the pathway level.

We have selected a number of pathways that warrant further investigation based on the fact that they were overrepresented among the previously identified altered genes. For these pathways, temporal gene regulatory network reconstruction was performed both within the mRNA level, as well as the effect of DNA copy number on mRNAs. Here we aimed to identify genes with the highest number of interactions within the pathway over time, which we called regulators as we believe they represent key altered genes for this pathway during cancer development. ID1 and PITX2 were identified as main regulators of the altered TGF-beta signaling pathway, BRWD3, and NF2 for mTOR signaling, while PIGT and DAPP1 for focal adhesion pathway. Functional validation experiments in the cell lines support the validity of our approach and the relevance of the identified genes.

This thesis sets the basis of integrative temporal statistical methodology in the direction of the differential expression analysis, gene regulatory network reconstruction

and identification of miRNA targets. The validity and applicability of our approaches were illustrated in the comprehensive investigation of molecular mechanisms underlying HPV-induced transformation in a cell line model. Our developed statistical methodology are important for an upcoming area in cancer research. Although currently in the literature multi-level time course experiments and corresponding datasets are rare, the expected introduction of blood-based tests for early cancer detection and treatment monitoring will generate molecular time course data in the near future.

Curriculum vitae

Viktorian Miok was born on October 27, 1985 in Zrenjanin, Serbia. In 2004 he graduated from high school Zrenjaninska Gimnazia in Zrenjanin. Viktorian studied Mathematics at the University of Belgrade, Serbia. He graduated in 2008, while in the same year he taught mathematics at the Economical High school in Zrenjanin. He received the Master of Science degree in Applied Statistics and Informatics from West University of Timisoara, Romania in 2010. During this study, he has also been employed as SAS Programmer at the Contact Research Organization Cmed in Timisoara. In the same year he has been given an opportunity to work as research fellow in statistics at the University of Bucharest, Romania under the supervision of Prof M Dumitrescu. In January 2012 he started his doctoral studies in statistics for integrative bioinformatics on joint project between the Department of Epidemiology & Biostatistics (Prof Dr MA van de Wiel) and Department of Pathology (Prof Dr P.J.F Snijders) of the Amsterdam University Medical Centers, which resulted in this thesis. Currently he is employed as Bioinformatician at Seven Bridges Genomics in Belgrade.