

Table of contents

1	Introduction	1
1.1	The importance of quantitative <i>in vivo</i> approach in biophysical studies . . .	1
1.2	Localization and mobility: two fundamental features of proteins in bacteria	4
1.3	Spatial organization of membrane receptor proteins is key to bacterial chemotaxis	6
1.4	Macromolecular crowding in biological systems	9
1.5	Bacterial nucleoids are not random polymers and live in the crowded cytoplasm	12
1.6	Single molecule fluorescence imaging approach allow for studying protein organization, mobility and interactions	14
1.7	Mother machine allows for nucleoid imaging in <i>E. coli</i> under physiological growth	16
1.8	Application of Super Resolution imaging to non-biological systems	18
1.9	Thesis outline and scope	19
2	Methods for bacterial chemoreceptor imaging	23
2.1	Introduction	23
2.1.1	PALM Imaging and Cluster-Size Analysis of Bacterial Chemoreceptors	27
2.1.2	LPA-SPT Analysis of Chemoreceptor Mobility	27
2.2	Materials	28
2.2.1	Strains and Plasmids	28
2.2.2	Growth Media	30
2.2.3	Buffer Solutions	30
2.2.4	Coverslip Cleaning	30
2.2.5	Cell Attachment and Drift Correction	31
2.2.6	Preparation of PALM Imaging Chamber	32
2.2.7	LPA-SPT Consumables for Flow Cell	32
2.2.8	General Microscopy	32
2.2.9	Additional Component for PALM	33

2.2.10	Additional Components for LPA-SPT	33
2.2.11	Fluidics	33
2.3	Methods	34
2.3.1	PALM Imaging of Bacterial Chemoreceptors	34
2.3.2	PALM Setup Summary	34
2.3.3	Chamber Preparation	36
2.3.4	Bacterial Cell Preparation	37
2.3.5	PALM Imaging Protocol	38
2.3.6	PALM data analysis	40
2.3.7	Cluster Analysis	42
2.3.8	Localized Photoactivation Single-Particle Tracking (LPA-SPT) of Chemoreceptors	43
2.3.9	Localized Photoactivation Path	44
2.3.10	LPA-SPT Sample Preparation	44
2.3.11	Cleaning of Flow Cell	46
2.3.12	Coverslip Mounting	46
2.3.13	Fluidic Switch Valve	47
2.3.14	LPA-SPT Imaging Protocol	47
2.3.15	LPA-SPT Data Analysis	49
2.3.16	Notes	51
2.4	Acknowledgments	54
3	Fluorescent toolkit for bacterial chromosome imaging	57
3.1	Introduction	58
3.2	Results	61
3.2.1	A novel fluorescent DNA label for <i>in vivo</i> chromosome imaging derived from the H-NS DNA binding domain (HNS-dbd).	61
3.2.2	High-resolution 2D and 3D imaging of nucleoids labeled via fluores- cent H-NS	65
3.2.3	Probing the mobility of DNA-associated proteins using photocon- vertible fluorophores.	67
3.2.4	Low-perturbation FROS system for studies on local gene compaction	71
3.2.5	A microfluidic chemostat (mother machine) allows for <i>in vivo</i> quan- titative measurements under physiological conditions	79
3.3	Discussion	82
3.4	Materials and Methods	86
3.4.1	Bacterial strains, plasmids and culture conditions	86

3.4.2	Microscopy	87
3.4.3	PALM and SPT data analysis	88
3.4.4	FROS data analysis	88
3.4.5	Mother Machine	89
3.5	Acknowledgments	93
4	Chemoreceptor cluster size and mobility under stimulation	95
4.1	Introduction	95
4.2	Results	98
4.2.1	PALM reveals extensive diversity in chemoreceptor spatial organization across cells	98
4.2.2	Cluster size is reduced upon attractant stimulation, and restored by adaptation	100
4.2.3	Global pattern of localization and local density of clusters are conserved under stimulation	102
4.2.4	Live-cell PALM confirms stimulus-induced destabilization of receptor clusters without chemical fixation	104
4.2.5	Direct observation of ligand-induced mobility changes by localized photoactivation single-particle tracking (LPA-SPT)	107
4.2.6	Stimulation by cognate ligands trigger an increase in the mobile fraction of chemoreceptors	110
4.2.7	Dynamics of cluster mobility under sustained stimulation reveals adaptation-dependent recovery	112
4.3	Discussion	114
4.4	Materials and methods	118
4.4.1	Bacterial strains and plasmids	118
4.4.2	Media and buffer	120
4.4.3	Cell culture	120
4.4.4	Sample preparation	121
4.4.5	Microscopy	123
4.4.6	Data analysis	125
4.5	Supporting Information	132
4.5.1	Molecular Counting with PALM	132
4.5.2	Functionality of the Tar-mEos2 fusion	140
4.5.3	Evidence that cognate ligand binding triggers changes in chemoreceptor cluster mobility	142
4.5.4	Dependence of the ligand-effect on chemoreceptor expression levels	146

4.5.5	Additional Supporting Figures	149
5	Nucleoid organization under increased crowding and confinement <i>in vivo</i>	153
5.1	Introduction	154
5.2	Results	157
5.2.1	Hyperosmotic shock in growing cells leads to transient crowding increase and nucleoid compaction	157
5.2.2	Hypoosmotic shock reduces crowding and increases nucleoid size .	160
5.2.3	Crowding plays an important role in setting the steady-state nucleoid size <i>in vivo</i>	163
5.2.4	Magnitude of nucleoid-size responses to hyperosmotic shock are comparable at different growth rates	164
5.2.5	Both crowding and nucleoid dynamics demonstrate hysteresis across hyperosmotic shock response and recovery	166
5.2.6	Nucleoid collapse, but not recovery, upon hyperosmotic shock follows a stereotyped trajectory	169
5.2.7	Metabolic shutdown dilates the nucleoid independently of crowding and inhibits recovery from collapse	170
5.3	Discussion	172
5.4	Materials and methods	179
5.4.1	Bacterial strains and plasmids.	179
5.4.2	Growth conditions	180
5.4.3	Overview of experimental setup and measurement strategy.	181
5.4.4	Microfluidic experiments in the mother machine	183
5.4.5	Microscopy	184
5.4.6	Quantitative data analysis and image processing.	185
5.5	Acknowledgments	187
	Appendix A Super resolution imaging of silicon nano-wires absorbtion profile	189
	Appendix References	202
	References	213